

# Categories for Dynamic Epistemic Logic

Kohei Kishida\*

Department of Computer Science  
University of Oxford  
Oxford, United Kingdom  
kohei.kishida@cs.ox.ac.uk

The primary goal of this paper is to recast the semantics of modal logic, and dynamic epistemic logic (DEL) in particular, in category-theoretic terms. We first review the category of relations and categories of Kripke frames, with particular emphasis on the duality between relations and adjoint homomorphisms. Using these categories, we then reformulate the semantics of DEL in a more categorical and algebraic form. Several virtues of the new formulation will be demonstrated: The DEL idea of updating a model into another is captured naturally by the categorical perspective—which emphasizes a family of objects and structural relationships among them, as opposed to a single object and structure on it. Also, the categorical semantics of DEL can be merged straightforwardly with a standard categorical semantics for first-order logic, providing a semantics for first-order DEL.

## 1 Introduction

*Dynamic epistemic logic* (DEL) is a powerful tool at the core of “logical dynamics” [8], a logical approach to the dynamics of information and interaction. Its semantics is general, flexible, and applicable to a wide range of informational processes in which rational agents update their knowledge and belief. It is also malleable and admits a variety of extra structures—e.g. probabilities, preferences, questions, awareness. It therefore forms a basis for logical studies of various aspects of agency in information and interaction.

The primary goal of this paper is to reformulate the standard semantics of DEL in category-theoretic terms.<sup>1</sup> One central idea of that semantics is that, to interpret DEL, we need to consider not just a single model but a family of models, in which one model is “updated” into another by a certain construction that models a given type of informational process. This is, in fact, a kind of idea that is treated naturally from the perspective of category theory. Category theory emphasizes a family of objects and structural relationships among them, as opposed to a single object and structure on it. Moreover, it can compare structural relationships at a “higher level” among different categories, e.g. between a category and another that is obtained by adding extra structure to the former. All this makes category theory excellent at capturing structural properties of a given family of models and constructions in a conceptually unifying fashion. And this paper will show that the semantics of DEL is an instance of this.

Section 2 will lay out Kripke semantics for propositional classical modal logic from a categorical perspective. Many of the concepts and facts covered in Section 2, such as subframes or duality results, are found in standard expositions such as [12, 10]; yet we will put more emphasis on the categorical structure of Kripke frames and on “higher” duality between relations and algebra operations. In Section 3 we will use the categorical structure of Kripke frames to shed new, categorical light on the standard semantics of

---

\*Kishida’s research has been supported by the grants FA9550-12-1-0136 of the U.S. AFOSR and EP/N018745/1 of EPSRC. An acknowledgment also goes to the anonymous referees for insightful comments and suggestions, which helped to improve the paper.

<sup>1</sup>See [3] for a clear and conceptual exposition of category theory.

DEL. We are not to propose a new semantics in this section, and the facts that will be covered are already known in literature (e.g. the standard exposition [16]). The point will instead be to use a categorical formulation and thereby to highlight structural properties in the standard semantics of DEL, uncovering the dual, algebraic ideas behind the semantics. Section 4 will give a demonstration of a virtue of our categorical, structural perspective, by showing how to extend DEL to the first order with a new, “sheaf” semantics.<sup>2</sup> Clearly, our knowledge and belief and their update often involve quantified propositions, and therefore can be subject to “first-order DEL”. It may nevertheless appear extremely complicated to introduce gadgets for quantification to the DEL framework. The structural approach, however, enables us to treat the DEL structure and the first-order structure as two modules to be simply combined. This will make obvious the conceptual power of the approach. Section 5 will discuss connections between our approach and some of the preceding categorical ones (such as the coalgebraic one). Then Section 6 will conclude the paper, referring to lines of future work.

This article adopts the following convention when displaying facts and results: Already known results are called “Facts”, with references attached in footnotes. Results that have not been explicitly stated before (to the best of the author’s knowledge) are called “Theorems” or “Corollaries” (the latter follow from already known results immediately).<sup>3</sup>

## 2 A Categorical Look at Kripke Semantics

This preliminary section lays out a categorical perspective on Kripke semantics for propositional classical modal logic. We mostly consider a single pair of unary modal operators  $\Box$  and  $\Diamond$ , but everything extends to a family of operator pairs (as we will see in the final paragraph of Subsection 2.3).

### 2.1 The Category of Relations

Let us first review basic facts about the category of binary relations. Given sets  $X$  and  $Y$ , we write  $R : X \rightarrow Y$  to mean that  $R$  is a relation “from  $X$  to  $Y$ ”, i.e.  $R \subseteq X \times Y$ . Relations  $R_1 : X \rightarrow Y$  and  $R_2 : Y \rightarrow Z$ , sharing the same  $Y$ , can be composed to form another  $R_1;R_2 : X \rightarrow Z$ , by defining  $wR_1;R_2u$  iff  $wR_1vR_2u$  for some  $v \in Y$ . The composition is also written  $R_2 \circ R_1$  (note the opposite orders of writing  $R_1$  and  $R_2$ ). The identity relation  $w = v$  on  $X$ , written  $1_X$ , is the identity of this composition, meaning that  $1_X;R = R = R;1_Y$  for every  $R : X \rightarrow Y$ . Then sets and binary relations form a category, **Rel**. This category comes with some extra structures, of which the most relevant to this article are the following:<sup>4</sup>

- **Rel** is a “dagger category”: Each relation  $R : X \rightarrow Y$  has its opposite,  $R^\dagger : Y \rightarrow X$ , so that  $vR^\dagger w$  iff  $wRv$ . This operation  $-^\dagger$  satisfies  $R^{\dagger\dagger} = R$  and  $(R_2 \circ R_1)^\dagger = R_1^\dagger \circ R_2^\dagger$ , and extends to a self-dual functor  $-^\dagger : \mathbf{Rel}^{\text{op}} \rightarrow \mathbf{Rel}$  by setting  $X^\dagger = X$  for each set  $X$ .<sup>5</sup>
- **Rel** is “locally posetal”: For each pair of sets  $X$  and  $Y$ , the set  $\mathbf{Rel}(X, Y)$  of relations from  $X$  to  $Y$  is a poset ordered by  $\subseteq$ . That is, relations  $R_1, R_2 : X \rightarrow Y$  satisfy the “higher” relation  $R_1 \subseteq R_2$  if

<sup>2</sup>A first-order extension of dynamic logic was given in [20]. The first attempt to extend DEL to the first order was [29], which introduced terms that referred to epistemic agents (and hence had a different format of logic than in this paper). Both of these extensions used constant domains for interpreting quantification. Constant domains can be seen as a (rather rigid) subcase of sheaves (and less flexible than sheaves in general); their axiomatization requires the so-called Barcan formula and other axioms be added to the simple union of modal logic and first-order logic (see Fact 5).

<sup>3</sup>We thank an anonymous reviewer for their suggestion of this convention.

<sup>4</sup>Categories with the following structures are studied e.g. in [33], where they are called “ordered categories with involution”.

<sup>5</sup>**Rel** admits an even stronger structure of “dagger compact (closed) category”, but this structure does not play an explicit rôle in this article. See Subsection 3.4.2 of [15] for **Rel** as a dagger compact category.

$wR_1v$  implies  $wR_2v$ . Moreover, posets  $\mathbf{Rel}(X, Y)$  and  $\mathbf{Rel}(Y, Z)$  interact with each other in such a way that if  $R_1 \subseteq R_2$  in  $\mathbf{Rel}(X, Y)$  and  $R_3 \subseteq R_4$  in  $\mathbf{Rel}(Y, Z)$  then  $R_3 \circ R_1 \subseteq R_4 \circ R_2$ .

- The two structures then interact in such a way that the functor  $\dagger$  gives order isomorphisms  $\dagger : \mathbf{Rel}(X, Y) \rightarrow \mathbf{Rel}(Y, X)$ ; i.e.,  $R_1 \subseteq R_2$  iff  $R_1^\dagger \subseteq R_2^\dagger$ .

The locally posetal structure makes  $\mathbf{Rel}$  a higher category with objects (“0-cells”)  $X$ , arrows (“1-cells”)  $R$  between objects, and higher arrows (“2-cells”)  $\subseteq$  between arrows (1-cells).<sup>6</sup> In addition,  $\mathbf{Rel}$  satisfies

1. the “law of modularity”:  $((R_2 \circ R_1) \cap R_3) \subseteq R_2 \circ (R_1 \cap (R_2^\dagger \circ R_3))$  (i.e., if  $wR_1; R_2u$  and  $wR_3u$  then there is  $v$  such that  $wR_1v$  and  $vR_2u$  and hence  $wR_3; R_2^\dagger v$ ).

Many properties of relations can be expressed with  $\subseteq$ . E.g.,  $R : X \rightarrow X$  is reflexive, i.e.  $w = v$  implies  $wRv$ , iff  $1_X \subseteq R$ . In particular, a relation  $R : X \rightarrow Y$  is a function iff both  $1_X \subseteq R^\dagger \circ R$  and  $R \circ R^\dagger \subseteq 1_Y$ . In addition, a function  $f : X \rightarrow Y$  is injective iff  $f^\dagger \circ f = 1_X$  and surjective iff  $f \circ f^\dagger = 1_Y$ . Functions are thus a subcase of relations. Moreover, the composition  $R_2 \circ R_1$  of relations is just the usual composition of functions when  $R_1$  and  $R_2$  are functions. So the category  $\mathbf{Sets}$  of sets and functions is a subcategory of  $\mathbf{Rel}$ . On the other hand,  $\mathbf{Sets}$  gives rise to  $\mathbf{Rel}$  as follows. A pair of functions  $f : Z \rightarrow X$  and  $g : Z \rightarrow Y$  from the same domain  $Z$  is called “jointly monic” if  $\langle f, g \rangle : Z \rightarrow X \times Y :: u \mapsto (f(u), g(u))$  is injective (or, equivalently,  $(f^\dagger \circ f) \cap (g^\dagger \circ g) = 1_Z$ ). Then a relation  $R : X \rightarrow Y$  corresponds to a jointly monic pair of functions, viz. the projections  $r_1 : R \rightarrow X :: (w, v) \mapsto w$  and  $r_2 : R \rightarrow Y :: (w, v) \mapsto v$  from the set  $R \subseteq X \times Y$ , so that the pair  $(r_1, r_2)$  “tabulates” the relation  $R : X \rightarrow Y$ , meaning that  $R = r_2 \circ r_1^\dagger$ .<sup>7</sup>

## 2.2 Relation-Modality Biduality

Kripke semantics uses binary relations to interpret unary modal operators. A Kripke frame is a set  $X$  paired with a binary relation  $R : X \rightarrow X$ , and a Kripke model is a Kripke frame  $(X, R)$  equipped with an assignment  $\llbracket - \rrbracket$  of subsets  $\llbracket p \rrbracket \subseteq X$  to propositional variables  $p$ . In fact we extend the notation to all propositions  $\varphi$ , so that  $w \in \llbracket \varphi \rrbracket \subseteq X$  means that  $\varphi$  is true at  $w$ . Now, given a relation  $R : X \rightarrow Y$ , define two monotone maps  $\exists_R, \forall_R : \mathcal{P}X \rightarrow \mathcal{P}Y$  by

$$\begin{aligned}\exists_R(S) &= \{v \in Y \mid w \in S \text{ for some } w \in X \text{ such that } wRv\}, \\ \forall_R(S) &= \{v \in Y \mid w \in S \text{ for all } w \in X \text{ such that } wRv\}.\end{aligned}$$

Then, for a relation  $R : X \rightarrow X$  on a set  $X$ ,  $\exists_{R^\dagger}, \forall_{R^\dagger} : \mathcal{P}X \rightarrow \mathcal{P}X$  interpret the “possibility” operator  $\diamond$  and the “necessity” operator  $\square$ , respectively—i.e.

$$\llbracket \diamond \varphi \rrbracket = \exists_{R^\dagger} \llbracket \varphi \rrbracket, \quad \llbracket \square \varphi \rrbracket = \forall_{R^\dagger} \llbracket \varphi \rrbracket. \quad (2)$$

An important property of  $\exists_-$  and  $\forall_-$  is that every relation  $R$  gives an adjunction (or “Galois connection”)  $\exists_R \dashv \forall_{R^\dagger}$ , meaning that  $\exists_R(S_1) \subseteq S_2$  iff  $S_1 \subseteq \forall_{R^\dagger}(S_2)$ . (And it also gives  $\exists_{R^\dagger} \dashv \forall_R$  via  $R^\dagger$ .) Therefore left adjoints  $\exists_R$  preserve arbitrary joins and right adjoints  $\forall_R$  preserve arbitrary meets. It also needs noting that a relation  $f : X \rightarrow Y$  is a function iff  $\exists_{f^\dagger} = \forall_{f^\dagger}$ , in which case  $\exists_{f^\dagger} = \forall_{f^\dagger}$  is the inverse-image map

<sup>6</sup>See Chapter XII of [36] for this type of higher categories, “bicategories”.  $\mathbf{Rel}$  appears in Subsection 1.5 (i) of [32] as an example of bicategory. A more general account of bicategories of relations is found in [11]. A similar approach, in terms of categories called “allegories”, is taken in Chapter 2 of [17], which also gives a thorough account of ideas in this subsection.

<sup>7</sup>The correspondence mentioned here is not quite 1–1. For two jointly monic pairs of functions  $(r_1 : Z \rightarrow X, r_2 : Z \rightarrow Y)$  and  $(r'_1 : Z' \rightarrow X, r'_2 : Z' \rightarrow Y)$ , if there is a bijection  $f : Z \rightarrow Z'$  such that  $r_i = r'_i \circ f$  for  $i = 1, 2$ , then the two pairs correspond to the same relation  $R \subseteq X \times Y$ . One can of course identify such isomorphic pairs of jointly monic pairs and force the correspondence to be 1–1.

$f^{-1} : \mathcal{P}Y \rightarrow \mathcal{P}X$ . So, for every function  $f$ , the map  $f^{-1} = \exists_{f^\dagger} = \forall_{f^\dagger}$  preserves all joins and meets, and moreover  $\exists_f \dashv f^{-1} \dashv \forall_f$ , which is one of the fundamental facts for categorical logic.<sup>8</sup>

One of the most fundamental categorical facts to the interpretation (2) is the equivalence of **Rel** and categories of complete atomic Boolean algebras (CABAs). Let  $\mathbf{CABA}_\vee$  and  $\mathbf{CABA}_\wedge$  be the categories of CABAs with all-join-preserving maps and with all-meet-preserving maps, respectively, and then

**Fact 1.**  $\exists_- :: R \mapsto \exists_R$  and  $\forall_- :: R \mapsto \forall_R$  extend to equivalences of categories  $\exists_- : \mathbf{Rel} \rightarrow \mathbf{CABA}_\vee$  and  $\forall_- : \mathbf{Rel} \rightarrow \mathbf{CABA}_\wedge$ , both sending a set  $X$  to its powerset  $\mathcal{P}X$ , while every CABA has the form  $\mathcal{P}X$ .<sup>9</sup> Putting this in “concrete” terms,

- The relations  $R : X \rightarrow Y$  correspond 1-1 to the all-join-preserving maps  $\exists_R : \mathcal{P}X \rightarrow \mathcal{P}Y$ , and also 1-1 to the all-meet-preserving maps  $\forall_R : \mathcal{P}X \rightarrow \mathcal{P}Y$ . In other words, for every pair of sets  $X$  and  $Y$ , each of  $\exists_-$  and  $\forall_-$  induces a bijection from  $\mathbf{Rel}(X, Y)$  to the set  $\mathbf{C}(\mathcal{P}X, \mathcal{P}Y)$  of arrows of  $\mathbf{C} = \mathbf{CABA}_\vee, \mathbf{CABA}_\wedge$  from  $\mathcal{P}X$  to  $\mathcal{P}Y$ .

In fact, higher versions of Fact 1 are relevant to modal logic. Recall that **Rel** is equipped with higher arrows between arrows, i.e. the relations  $\subseteq$  among relations  $R_1, R_2 : X \rightarrow Y$ . Similarly,  $\mathbf{C} = \mathbf{CABA}_\vee, \mathbf{CABA}_\wedge$  are also equipped with the relation  $\leq$  among arrows  $h_1, h_2 : \mathcal{P}X \rightarrow \mathcal{P}Y$ , by setting  $h_1 \leq h_2$  iff  $h_1(S) \subseteq h_2(S)$  for all  $S \in \mathcal{P}X$ , making  $\mathbf{C}(\mathcal{P}X, \mathcal{P}Y)$  a poset. Then

**Fact 2.**  $\exists_- : \mathbf{Rel} \rightarrow \mathbf{CABA}_\vee$  is a (higher) equivalence,<sup>10</sup> meaning that

- $R_1 \subseteq R_2$  iff  $\exists_{R_1} \leq \exists_{R_2}$ ,

i.e., each bijection  $\exists_- : \mathbf{Rel}(X, Y) \rightarrow \mathbf{CABA}_\vee(\mathcal{P}X, \mathcal{P}Y)$  is an order isomorphism.

Yet there are more versions of this result that are less frequently mentioned but equally important. Since **Rel** has two levels of arrows,  $R$  and  $\subseteq$ , there are four types of (higher) functors  $F$  from **Rel** to another higher category  $\mathbf{C}$ , viz.,  $F : \mathbf{Rel} \rightarrow \mathbf{C}$ , with the direction of neither  $R$  nor  $\subseteq$  flipped;  $F : \mathbf{Rel}^{\text{op}} \rightarrow \mathbf{C}$ , with just  $R$  flipped;  $F : \mathbf{Rel}^{\text{co}} \rightarrow \mathbf{C}$ , with just  $\subseteq$  flipped;  $F : \mathbf{Rel}^{\text{coop}} \rightarrow \mathbf{C}$ , with both  $R$  and  $\subseteq$  flipped.<sup>11</sup> Therefore there can be four versions of equivalence (or duality).

**Corollary 1.**  $\forall_- : \mathbf{Rel}^{\text{co}} \rightarrow \mathbf{CABA}_\wedge$  is a “2-cell duality”, i.e. an equivalence flipping  $\subseteq$  (but not  $R$ ). Concretely put,

- $R_1 \subseteq R_2$  iff  $\forall_{R_2} \leq \forall_{R_1}$  (note the reversed order),

i.e., each bijection  $\forall_- : \mathbf{Rel}(X, Y) \rightarrow \mathbf{CABA}_\wedge(\mathcal{P}X, \mathcal{P}Y)$  is an order-reversing isomorphism.

Moreover, composing  $\exists_-$  and  $\forall_-$  with the self-dual functor  $\dagger : \mathbf{Rel}^{\text{op}} \rightarrow \mathbf{Rel}$ , which is a “1-cell duality”, i.e. an equivalence flipping  $R$  but not  $\subseteq$ , we obtain

**Corollary 2.**  $\exists_{-\dagger} : \mathbf{Rel}^{\text{op}} \rightarrow \mathbf{CABA}_\vee$  is a 1-cell duality, and  $\forall_{-\dagger} : \mathbf{Rel}^{\text{coop}} \rightarrow \mathbf{CABA}_\wedge$  is a “biduality”, i.e. an equivalence flipping both  $R$  and  $\subseteq$ . Concretely put,

3.  $\exists_{(R_2 \circ R_1)^\dagger} = \exists_{R_1^\dagger} \circ \exists_{R_2^\dagger}$  and  $\forall_{(R_2 \circ R_1)^\dagger} = \forall_{R_1^\dagger} \circ \forall_{R_2^\dagger}$  (note the flipped orders of composition),
4.  $R_1 \subseteq R_2$  iff  $\exists_{R_1^\dagger} \leq \exists_{R_2^\dagger}$ ,

<sup>8</sup>The idea that  $\square$  and  $\diamond$  are a relational generalization of  $\forall_f$  and  $\exists_f$  is laid out in [22] from a more general categorical perspective of “bicategories of relations and spans”. As observed in [22], we can define modal operators  $\blacklozenge$  and  $\blacksquare$  that are “opposite” to  $\diamond$  and  $\square$ , and interpret them with  $\exists_R$  and  $\forall_R$ ; then we have adjunctions  $\blacklozenge \dashv \square$  and  $\diamond \dashv \blacksquare$ , which also appear in [31]. These adjunctions are typical of the “past” and “future” modalities of temporal logic, as observed in [25].

<sup>9</sup>See, e.g., Exercise 5.2.5 in [23] for essentially the same fact.

<sup>10</sup>See Subsection 1.5 (i) of [32] for essentially the same fact.

<sup>11</sup> $\mathbf{Rel}^{\text{op}}$ ,  $\mathbf{Rel}^{\text{co}}$ , and  $\mathbf{Rel}^{\text{coop}}$  are **Rel** with just  $R$  flipped; with just  $\subseteq$  flipped; and with both  $R$  and  $\subseteq$  flipped. See Subsection 1.6 of [32].

5.  $R_1 \subseteq R_2$  iff  $\forall_{R_2^\dagger} \leq \forall_{R_1^\dagger}$  (note the reversed order),

i.e.,  $\exists_{-\dagger}$  and  $\forall_{-\dagger}$  induce order-preserving isomorphisms  $\exists_{-\dagger} : \mathbf{Rel}(Y, X) \rightarrow \mathbf{CABA}_\vee(\mathcal{P}X, \mathcal{P}Y)$  and order-reversing isomorphisms  $\forall_{-\dagger} : \mathbf{Rel}(Y, X) \rightarrow \mathbf{CABA}_\wedge(\mathcal{P}X, \mathcal{P}Y)$ , respectively.

Thus, (2) means that the modal operators  $\diamond$  and  $\square$  are duals to the relations  $R : X \rightarrow X$ , in such a “higher” way that the relation  $\subseteq$  among the latter corresponds to the relation  $\leq$  among the former (e.g. the reflexivity of  $R$ , i.e.  $1_X \subseteq R$ , is equivalent by (4) to  $1_{\mathcal{P}X} \leq \exists_{R^\dagger}$  and by (5) to  $\forall_{R^\dagger} \leq 1_{\mathcal{P}X}$ , i.e.  $\varphi \vdash \diamond\varphi$  and  $\square\varphi \vdash \varphi$ ).<sup>12</sup> This higher duality plays a fundamental rôle in this article as well as in Kripke semantics in general.

One more fact that will prove useful is the “Beck-Chevalley condition”:

**Corollary 3.** If the following diagram is a pullback in **Sets**, then  $p \circ q^\dagger = f^\dagger \circ g$ .<sup>13</sup>

$$\begin{array}{ccc}
 Y \times_X Z & \xrightarrow{q} & Z \\
 \downarrow p & \lrcorner & \downarrow g \\
 Y & \xrightarrow{f} & X
 \end{array} \tag{6}$$

### 2.3 Categories of Kripke Frames

Let us now consider categories of Kripke frames. A *monotone* map from a Kripke frame  $(X, R_X)$  to another  $(Y, R_Y)$  is a function  $f : X \rightarrow Y$  that preserves relation, i.e. such that  $wR_X v$  implies  $f(w)R_Y f(v)$ . Observe that this can equivalently be written as either of the following.

7.  $R_X \subseteq f^\dagger \circ R_Y \circ f$  (i.e.,  $wR_X v$  implies  $wf w'R_Y v' f^\dagger v$  for some  $w', v' \in Y$ ),

8.  $f \circ R_X \subseteq R_Y \circ f$  (i.e.,  $wR_X v f v'$  implies  $wf w'R_Y v'$  for some  $w' \in Y$ ).

The formulation (8) strengthens to  $f$  being a *bounded morphism*, i.e. satisfying both (8) and

9.  $R_Y \circ f \subseteq f \circ R_X$  (i.e.,  $wf w'R_Y v'$  implies  $wR_X v f v'$  for some  $v \in X$ ),

i.e. satisfying

10.  $f \circ R_X = R_Y \circ f$ .

Let us write **Kr** for the category of Kripke frames and monotone maps, and **Kr<sub>B</sub>** for its subcategory of bounded morphisms.

The duality observed in Subsection 2.2 immediately entails duality results between Kripke frames and “CABAs with operators” (CABAOs), i.e. CABAs equipped with all-join-preserving operators  $\diamond$ . The isomorphisms  $\exists_{-\dagger} : \mathbf{Rel}(X, X) \rightarrow \mathbf{CABA}_\vee(\mathcal{P}X, \mathcal{P}X)$  in Corollary 2 mean that the Kripke frames  $(X, R)$  correspond 1-1 to the CABAOs  $(\mathcal{P}X, \diamond)$ . Moreover, while the functions  $f : X \rightarrow Y$  and the CABA homomorphisms  $h : \mathcal{P}Y \rightarrow \mathcal{P}X$  are dual to each other, Corollary 2 further implies (by  $f^{-1} = \exists_{f^\dagger} = \forall_{f^\dagger}$ ) that (10) is equivalent to

11.  $\exists_{R_X^\dagger} \circ f^{-1} = f^{-1} \circ \exists_{R_Y^\dagger}$  (or equivalently  $\forall_{R_X^\dagger} \circ f^{-1} = f^{-1} \circ \forall_{R_Y^\dagger}$ ),

<sup>12</sup>See [27] for more on correspondence results via the higher dualities.

<sup>13</sup>**Sets** satisfies the Beck-Chevalley condition, meaning that the pullback entails  $\exists_q \circ p^{-1} = g^{-1} \circ \exists_f$ . See Section IV.9 of [37]. This implies (6) by Corollary 2.

i.e.,  $f^{-1}$  being a CBAO homomorphism, i.e. a CABA homomorphism that moreover preserves  $\diamond$  (and  $\square$ ), from  $(\mathcal{P}Y, \exists_{R_Y^\dagger})$  to  $(\mathcal{P}X, \exists_{R_X^\dagger})$ . Therefore the category  $\mathbf{Kr}_B$  is dual to the category **CBAO** of CBAOs and CBAO homomorphisms.<sup>14</sup> In fact, let us call a CABA homomorphism  $h$  “continuous” if it has  $\diamond \circ h \leq h \circ \diamond$ , and then Corollary 2 implies that (7)–(8) are equivalent to

12.  $\exists_{R_X^\dagger} \leq f^{-1} \circ \exists_{R_Y^\dagger} \circ \exists_f$ , or equivalently  $\exists_{R_X^\dagger} \circ f^{-1} \leq f^{-1} \circ \exists_{R_Y^\dagger}$  (or  $f^{-1} \circ \forall_{R_Y^\dagger} \circ \forall_f \leq \forall_{R_X^\dagger}$  or  $f^{-1} \circ \forall_{R_Y^\dagger} \leq \forall_{R_X^\dagger} \circ f^{-1}$ ),

i.e. the continuity of  $f^{-1}$ . Hence the category  $\mathbf{Kr}$  is dual to the category **CBAO<sub>C</sub>** of CBAOs and continuous CABA homomorphisms [19]. We should stress, however, that these duality results are merely derivative, and that the dualities in Subsection 2.2 are more fundamental. It is the latter duality that we will take essential advantage of throughout this article.

We have so far considered a single pair of operators  $\square$  and  $\diamond$ , but in epistemic logic we often take a set  $A$  of agents and consider a pair of operators  $[\alpha]$  (also written  $K_\alpha$ , for “ $\alpha$  knows that”) and  $\langle \alpha \rangle$  for each agent  $\alpha \in A$ . To interpret this  $A$ -indexed set of operator pairs, a Kripke frame  $X$  needs to be equipped with an  $A$ -indexed set of relations  $R_\alpha : X \rightarrow X$  as well. Let us say that a function  $f : X \rightarrow Y$  from a Kripke frame  $(X, R_X^\alpha)_{\alpha \in A}$  to another  $(Y, R_Y^\alpha)_{\alpha \in A}$  is monotone if it preserves every  $R_X^\alpha$  by satisfying (7)–(8) (with  $R_X^\alpha$  in place of  $R_X$ ), and a bounded morphism if it satisfies (7)–(10) for every  $R_X^\alpha$  (in place of  $R_X$ ). Then the Kripke frames with  $A$ -many relations and their monotone maps or bounded morphisms form categories  $\mathbf{Kr}_A$  and  $\mathbf{Kr}_{B,A}$ , subsuming  $\mathbf{Kr}$  and  $\mathbf{Kr}_B$  above as just a special case with  $A$  a singleton. The duality results in this section carry over straightforwardly to  $\mathbf{Kr}_A$  and  $\mathbf{Kr}_{B,A}$ , with respect to CABAs with  $A$ -many operators.

## 2.4 Topological Constructions for Kripke Frames

Having introduced two categories of Kripke frames, it may appear to be a natural question which of the two we should adopt as “the” category of Kripke frames. The answer is, however, that we need both  $\mathbf{Kr}$  and  $\mathbf{Kr}_B$ . The significance of  $\mathbf{Kr}_B$  is fairly obvious and well studied. Bounded morphisms are dual to homomorphisms preserving  $\diamond$  and  $\square$  as well as all the other connectives, and therefore closely connected to the preservation of modal logic. Indeed, the bisimulations are precisely the “relations in  $\mathbf{Kr}_B$ ” (see the final paragraph of Subsection 3.1). By the same token, in the coalgebraic approach to Kripke semantics, the kind of homomorphisms considered are those corresponding to bounded morphisms, and hence the considered category of coalgebras is equivalent to  $\mathbf{Kr}_B$  (see Section 5 for more on the connection to the coalgebraic approach). Quite arguably,  $\mathbf{Kr}_B$  plays a more prominent rôle than  $\mathbf{Kr}$  does, as long as the “static” modal logic is concerned. Nevertheless, this statement no longer applies to the semantics of dynamic epistemic logic (DEL). Many of the semantic constructions crucial for DEL take place in  $\mathbf{Kr}$  but not in  $\mathbf{Kr}_B$ . Indeed, to let DEL show interesting behaviors, it is essential to use monotone maps and not bounded morphisms.

The category  $\mathbf{Kr}$  admits a wide range of constructions that are directly connected to ones in **Sets** using sets and functions. They are due to

**Fact 3.**  $\mathbf{Kr}$  is “topological over **Sets**”,<sup>15</sup> meaning, concretely, the following. Given any family of functions  $f_i : X \rightarrow Y_i$  ( $i \in I$ ) to Kripke frames  $(Y_i, R_i)$ , the relation

$$wR_X v \iff f_i(w)R_i f_i(v) \text{ for all } i \in I, \quad \text{i.e.} \quad R_X = \bigcap_{i \in I} (f_i^\dagger \circ R_i \circ f_i),$$

<sup>14</sup>This duality was first shown in [40]. See also [10].

<sup>15</sup>See Section 21 of [1] for the definition and nice properties of topological categories. (It may need noting that [1] refers to  $\mathbf{Kr}$  as **Rel**.) This subsection refers to Definitions 21.1 and 21.7, Example 21.8, Propositions 21.30 and 21.31, Theorem 21.9, and Proposition 21.15.

is the (unique) “initial lift” of  $\{f_i\}_{i \in I}$ , i.e. the relation on  $X$  such that, given any function  $g : Z \rightarrow X$ , all  $f_i \circ g$  are monotone from a frame  $(Z, R_Z)$  iff  $g$  is.

(In fact, Fact 3 holds of  $\mathbf{Kr}_A$  in general, again with  $R_X^\alpha$  in place of  $R_X$ .) One may note that the relation  $R_X$  in Fact 3 is the largest relation on  $X$  preserved by all  $f_i$ , since, for every relation  $R$  on  $X$ ,

$$R \subseteq R_X \iff R \subseteq f_i^\dagger \circ R_i \circ f_i \text{ (i.e. } f_i \text{ preserves } R) \text{ for all } i \in I. \quad (13)$$

It is easy to observe that initial lifts preserve many properties of relations such as reflexivity, transitivity, and symmetry. Then the full subcategories of  $\mathbf{Kr}$  given by those properties and combinations thereof, such as **Preord** of the preorders (i.e. reflexive and transitive relations) and **Equiv** of the equivalence relations, are said to be “initially closed”. It follows that these subcategories are also topological over **Sets**, and that the inclusion functors have left-adjoints.<sup>16</sup> E.g., the left adjoint  $F : \mathbf{Kr} \rightarrow \mathbf{Preord}$  sends a Kripke frame  $(X, R)$  to  $(X, R^*)$ , where  $R^*$  is the reflexive and transitive closure of  $R$ .

One consequence of  $\mathbf{Kr}$ , or a subcategory such as **Preord**, being topological over **Sets** is that it also has “final lifts”, dual to initial lifts of Fact 3. E.g., given a family of preorders  $(X, R_\alpha)$  ( $\alpha \in A$ ) on the same set  $X$ , such as “epistemic” relations  $R_\alpha$  of agents  $\alpha \in A$ , consider an  $A$ -indexed family of identity maps  $\{1_X\}_{\alpha \in A}$  in **Sets**; then its final lift in **Preord** comes with the epistemic relation for the “common knowledge” of the group  $A$ , i.e.  $(\bigcup_\alpha R_\alpha)^*$ .<sup>17</sup>

Another consequence, more relevant to this article, is that the forgetful functor  $U : \mathbf{Kr} \rightarrow \mathbf{Sets}$  to the complete and cocomplete category **Sets** lifts limits and colimits—meaning that, given any (small) diagram  $D$  in  $\mathbf{Kr}$ , its (co)limit exists on the (co)limit of  $U \circ D$  in **Sets**. Most notably,

14. Given a family of Kripke frames  $(Y_i, R_i)$  ( $i \in I$ ), its product in  $\mathbf{Kr}$ ,  $(X, R_X)$ , is defined on the cartesian product  $X = \prod_{i \in I} Y_i$  by taking  $R_X = \bigcap_{i \in I} (p_i^\dagger \circ R_i \circ p_i)$  for the projections  $p_i : X \rightarrow Y_i$ .
15. Let  $i : S \hookrightarrow X$  be an inclusion map. Then  $(S, R_S)$  is a subframe of a Kripke frame  $(X, R_X)$ , i.e.  $R_S = i^\dagger \circ R_X \circ i$ , iff  $i$  is a regular mono from  $(S, R_S)$  to  $(X, R_X)$  in  $\mathbf{Kr}$ .

These constructions, and their canonical maps  $p_i$  and  $i$ , are crucial to the semantics of DEL, as we will see in Section 3. Pullbacks in  $\mathbf{Kr}$  will also play a key rôle later in Subsection 4.3. In particular, observe

**Theorem 1.** *The pullback of a bounded morphism in  $\mathbf{Kr}$  is a bounded morphism.*<sup>18</sup>

*Proof.* Let (6) be a pullback in  $\mathbf{Kr}$ , let  $R_X$  be the relation on  $X$ , similarly for  $Y, Z$ , and  $Y \times_X Z$ , and let  $g$  be a bounded morphism. Then  $p$  satisfies (9) as follows, by (7) for  $f$ ; the commuting of (6); (10) for  $g$ ; Corollary 3; the law of modularity (1); and the definition of  $R_{Y \times_X Z}$  as the initial lift of  $p$  and  $g$ .

$$\begin{aligned} R_Y \circ p &\subseteq (R_Y \circ p) \cap (f^\dagger \circ R_X \circ f \circ p) = (R_Y \circ p) \cap (f^\dagger \circ R_X \circ g \circ q) \\ &= (R_Y \circ p) \cap (f^\dagger \circ g \circ R_Z \circ q) \\ &= (R_Y \circ p) \cap (p \circ q^\dagger \circ R_Z \circ q) \\ &\subseteq p \circ ((p^\dagger \circ R_Y \circ p) \cap (q^\dagger \circ R_Z \circ q)) = p \circ R_{Y \times_X Z}. \quad \square \end{aligned}$$

It needs stressing, however, that the canonical maps of “topological” constructions in this subsection are not in general bounded morphisms, and hence do not live in  $\mathbf{Kr}_B$ . Indeed, as we will see, they must not be bounded morphisms for DEL to show interesting behaviors.

<sup>16</sup>There are properties that are not preserved by initial lifts. E.g., antisymmetry is not; in fact, the category of posets is not topological over **Sets**.

<sup>17</sup>See Section 2.3 of [16], as well as [6, 7], for common knowledge. We do not treat its logic in this article.

<sup>18</sup>This is a straightforward analogue of the already known fact that, in the category of topological spaces, the pullback of an open map is open. See Proposition 1 in Section V.4 of [24].

### 3 A Categorical Look at Dynamic Epistemic Logic

This section shows how to use the categorical structure of Section 2 to reformulate the standard semantics of dynamic epistemic logic (DEL) structurally. We will first review the simpler subcase of *public announcement logic* (PAL) in Subsection 3.1, and then expand it to the general DEL in Subsection 3.2.

#### 3.1 Public Announcement Logic

Regular monos  $i$  of  $\mathbf{K}r$  in (15) are used to interpret PAL. This logic has unary operators  $[\sigma!]$  and  $\langle\sigma!\rangle$  for all of its propositions  $\sigma$ . The proposition  $[\sigma!]\varphi$  is intended to mean “ $\varphi$  will be the case after  $\sigma$  is publicly and truthfully announced (or observed)”, and interpreted as follows: Given a Kripke model  $(X, R_X, \llbracket - \rrbracket_X)$  and a subset  $S = \llbracket \sigma \rrbracket_X$  with inclusion  $i : S \hookrightarrow X$ , let  $(S, R_S, \llbracket - \rrbracket_S)$  be the submodel on  $S$ —which is defined by  $R_S = i^\dagger \circ R_X \circ i$  and  $\llbracket p \rrbracket_S = i^{-1} \llbracket p \rrbracket_X$  for atomic  $p$ . Then

16.  $w \in \llbracket [\sigma!]\varphi \rrbracket_X$  iff either  $w \notin \llbracket \sigma \rrbracket_X$  or  $w \in \llbracket \varphi \rrbracket_S$  (note the subscripts), i.e., iff  $v \in \llbracket \varphi \rrbracket_S$  for all  $v \in S$  such that  $viw$ . In short,  $\llbracket [\sigma!]\varphi \rrbracket_X = \forall_i \llbracket \varphi \rrbracket_S$ .

Similarly (or De Morgan-dually),

17.  $w \in \llbracket \langle\sigma!\rangle\varphi \rrbracket_X$  iff both  $w \in \llbracket \sigma \rrbracket_X$  and  $w \in \llbracket \varphi \rrbracket_S$ , i.e., iff  $v \in \llbracket \varphi \rrbracket_S$  for some  $v \in S$  such that  $viw$ . In short,  $\llbracket \langle\sigma!\rangle\varphi \rrbracket_X = \exists_i \llbracket \varphi \rrbracket_S$ .

One may contrast (16) and (17) to

$$\forall_i \circ i^{-1} \llbracket \varphi \rrbracket_X = \llbracket \sigma \Rightarrow \varphi \rrbracket_X, \quad \exists_i \circ i^{-1} \llbracket \varphi \rrbracket_X = \llbracket \sigma \wedge \varphi \rrbracket_X. \quad (18)$$

So, although generally  $\llbracket \varphi \rrbracket_S \neq i^{-1} \llbracket \varphi \rrbracket_X$ , for atomic  $p$  we have  $\llbracket p \rrbracket_S = i^{-1} \llbracket p \rrbracket_X$  by definition, and hence have a “reduction axiom”  $[\sigma!]p \equiv (\sigma \Rightarrow p)$  by

$$\llbracket [\sigma!]p \rrbracket_X = \forall_i \llbracket p \rrbracket_S = \forall_i \circ i^{-1} \llbracket p \rrbracket_X = \llbracket \sigma \Rightarrow p \rrbracket_X.$$

Reduction axioms, taken together for atomic sentences and for all the “static” connectives, completely axiomatize PAL by reducing it to the static modal logic. Proofs for reduction axioms for connectives are:

19. Because  $\forall_i$  preserves meets,

$$\llbracket [\sigma!](\varphi \wedge \psi) \rrbracket_X = \forall_i (\llbracket \varphi \rrbracket_S \cap \llbracket \psi \rrbracket_S) = \forall_i \llbracket \varphi \rrbracket_S \cap \forall_i \llbracket \psi \rrbracket_S = \llbracket [\sigma!]\varphi \wedge [\sigma!]\psi \rrbracket_X.$$

20. A CABA homomorphism,  $i^{-1}$  preserves  $\neg$ . And  $i^\dagger \circ i = 1_S$ , or dually  $i^{-1} \circ \forall_i = 1_{\mathcal{P}(S)}$ , since  $i$  is an injection. Therefore  $\neg_S = \neg_S \circ i^{-1} \circ \forall_i = i^{-1} \circ \neg_X \circ \forall_i$ . Hence

$$\llbracket [\sigma!]\neg\varphi \rrbracket_X = \forall_i \circ \neg_S \llbracket \varphi \rrbracket_S = \forall_i \circ i^{-1} \circ \neg_X \circ \forall_i \llbracket \varphi \rrbracket_S = \llbracket \sigma \Rightarrow \neg[\sigma!]\varphi \rrbracket_X.$$

21.  $R_S = i^\dagger \circ R_X \circ i$  dually means  $\forall_{R_S^\dagger} = i^{-1} \circ \forall_{R_X^\dagger} \circ \forall_i$ . Therefore

$$\llbracket [\sigma!]\Box\varphi \rrbracket_X = \forall_i \circ \forall_{R_S^\dagger} \llbracket \varphi \rrbracket_S = \forall_i \circ i^{-1} \circ \forall_{R_X^\dagger} \circ \forall_i \llbracket \varphi \rrbracket_S = \llbracket \sigma \Rightarrow \Box[\sigma!]\varphi \rrbracket_X.$$

These algebraic proofs are straightforward applications of properties of the duality  $\forall_{-\dagger}$ . In particular, it should be noted that (21), the reduction via  $\Box$ , is simply a dual to the equality of relations  $R_S \circ i^\dagger = i^\dagger \circ R_X \circ i \circ i^\dagger$ .

A perspective on (16)–(17) that has been guiding the study of the semantics of PAL, and indeed of DEL (see e.g. [6]), is that  $[\sigma!]$  and  $\langle\sigma!\rangle$  are interpreted by  $\forall_i$  and  $\exists_i$ , and therefore are the modal



operators of the relation  $i^\dagger$  (called a “transition relation” in [6]), similarly to  $\Box$  and  $\Diamond$  interpreted by  $\forall_{R^\dagger}$  and  $\exists_{R^\dagger}$  of  $R$  as in (2). One difference is that, whereas  $R$  is a relation on the same set,  $i^\dagger$  is between different sets. Thus PAL, and DEL in general, generalize Kripke semantics by using relations  $R : X \rightarrow Y$  between different Kripke frames to interpret modal operators. In studying this general setting, it proves helpful to use the relation-modality dualities of Subsection 2.2 (and not just the derivative dualities of Subsection 2.3 between Kripke frames and CBAOs). It may also be interesting to note that  $\sigma \Rightarrow -$  and  $\sigma \wedge -$  in (18), which play an essential rôle in reduction axioms, are modal operators, too, viz. those of the relation  $i \circ i^\dagger : X \rightarrow X$ . This is the reason the relation-modality duality  $\forall_{-^\dagger}$  is applicable in (21).

A point of caution here for our categorical approach is that, in general,  $R : X \rightarrow Y$  is neither a structure on a Kripke frame (an object of the category  $\mathbf{Kr}$ ) nor a monotone map (an arrow of  $\mathbf{Kr}$ ). So, to accommodate it in terms of  $\mathbf{Kr}$ , we use the idea of tabulation from Subsection 2.1: A relation  $R : X \rightarrow Y$  corresponds to the pair of projections  $r_1 : R \rightarrow X$  and  $r_2 : R \rightarrow Y$  from the set  $R \subseteq X \times Y$ , so that  $R = r_2 \circ r_1^\dagger$ . Indeed, given Kripke frames on  $X$  and  $Y$ , Fact 3 gives a canonical Kripke frame on  $R \subseteq X \times Y$  from which  $r_1$  and  $r_2$  are monotone. Then  $\forall_{R^\dagger} = \forall_{r_1} \circ r_2^{-1}$  and  $\exists_{R^\dagger} = \exists_{r_1} \circ r_2^{-1}$ ; hence  $\forall_{R^\dagger}$  and  $\exists_{R^\dagger}$  of all relations  $R$  can be obtained by  $\exists_f \dashv f^{-1} \dashv \forall_f$  of monotone maps  $f$ . This trick, using monotone maps  $r_1$  and  $r_2$  of  $\mathbf{Kr}$ , always works for any relation  $R \subseteq X \times Y$ . On the other hand, bounded morphisms of  $\mathbf{Kr}_B$  do not always work, since  $r_1$  and  $r_2$  are both bounded morphisms if and only if  $R$  is a bisimulation. (We will see an even more crucial rôle of  $\mathbf{Kr}$  at the end of Subsection 3.2.)

### 3.2 Dynamic Epistemic Logic

Let us now consider the Baltag-Moss-Solecki semantics of DEL [7] and observe how product update in it can be treated categorically. Take two Kripke frames,  $(X, R_X)$  and  $(E, R_E)$ , and regard the former as an “epistemic model” and the latter as an “event model”. So, let us assume that  $(X, R_X)$  is equipped with an interpretation  $\llbracket \text{Pre}(e) \rrbracket_X \subseteq X$  of the precondition  $\text{Pre}(e)$  of every event  $e \in E$  (or we can take a Kripke model  $(X, R_X, \llbracket - \rrbracket_X)$  on  $(X, R_X)$ ); we write  $i_e : \llbracket \text{Pre}(e) \rrbracket_X \hookrightarrow X$  for the inclusion maps. Then the product model of the two frames, obtained by “updating”  $(X, R_X)$  with  $(E, R_E)$ , is defined on the disjoint union of  $i_e$ , i.e. the subset

$$X \otimes E = \sum_{e \in E} \llbracket \text{Pre}(e) \rrbracket_X = \{(w, e) \in X \times E \mid w \in \llbracket \text{Pre}(e) \rrbracket_X\} \quad \text{of} \quad X \times E = \sum_{e \in E} X. \quad (22)$$

The “epistemic” relation  $R_{X \otimes E}$  on  $X \otimes E$  is defined as the subframe of the product  $(X \times E, R_{X \times E})$  of  $(X, R_X)$  and  $(E, R_E)$ , using (14) and (15). This amounts to

$$(w_1, e_1) R_{X \otimes E} (w_2, e_2) \iff w_1 R_X w_2 \text{ and } e_1 R_E e_2, \quad \text{i.e.,} \quad R_{X \otimes E} = (p_X^\dagger \circ R_X \circ p_X) \cap (p_E^\dagger \circ R_E \circ p_E)$$

for the projections  $p_X : X \otimes E \rightarrow X :: (w, e) \mapsto w$  and  $p_E : X \otimes E \rightarrow E :: (w, e) \mapsto e$ . In short, it is the initial lift of  $p_X$  and  $p_E$ . In addition, given a Kripke model  $\llbracket - \rrbracket_X$  on  $X$ , it induces an updated Kripke model on  $X \otimes E$  by  $\llbracket p \rrbracket_{X \otimes E} = p_X^{-1} \llbracket p \rrbracket_X$  for atomic  $p$ .<sup>19</sup>

Let us analyze this construction a bit further, using the following diagram (for each  $e \in E$ ).

$$\begin{array}{ccc} X \otimes E = \sum_{e' \in E} \llbracket \text{Pre}(e') \rrbracket_X & \xleftarrow{i} & \sum_{e' \in E} X = X \times E \xrightarrow{p'_E} E \\ \uparrow q_e \quad \lrcorner & & \uparrow q'_e \quad \downarrow p'_X \\ \llbracket \text{Pre}(e) \rrbracket_X & \xleftarrow{i_e} & X \end{array} \quad (23)$$

<sup>19</sup>This is the case without “factual change”. A version with factual change [9] can also be treated categorically.

Here  $p'_X$  and  $p'_E$  are the obvious projections, so that  $p_X = p'_X \circ i$  and  $p_E = p'_E \circ i$ . And  $q_e$  and  $q'_e$  are the “coproduct injections”  $w \mapsto (w, e)$ . The inclusion  $i : X \otimes E \hookrightarrow X \times E$  has  $i \circ q_e = q'_e \circ i_e$  (by its definition as  $i = \sum_{e' \in E} i_{e'}$ ), while  $p'_X \circ q'_e = 1_X$  (since  $p'_X$  equals the trivial “cotuple”  $[1_X]_{e' \in E}$ ), and therefore  $p_X \circ q_e = p'_X \circ i \circ q_e = p'_X \circ q'_e \circ i_e = i_e$ .

Given this construction, for each  $e \in E$  the canonical functions  $i_e$  and  $q_e$  tabulate a relation  $R_e = q_e \circ i_e^\dagger : X \rightarrow X \otimes E$ ; i.e.,  $wR_e(w', e')$  iff  $w = w' \in \llbracket \text{Pre}(e) \rrbracket_X$  and  $e = e'$ , or  $wR_e v$  iff  $vp_X w$  and  $vp_E e$ . Corollary 3 implies  $R_e = q_e \circ i_e^\dagger = i_e^\dagger \circ q'_e$  since the square in (23) is a pullback. This relation, and its duals  $\forall_{R_e^\dagger} = \forall_{i_e} \circ q_e^{-1}$  and  $\exists_{R_e^\dagger} = \exists_{i_e} \circ q_e^{-1}$ , are then used to interpret the dynamic operators  $[E, e]$  and  $\langle E, e \rangle$ ; the proposition  $[E, e]\varphi$  is supposed to mean “ $\varphi$  will be the case after the event  $e$  takes place”. The interpretation, similar to (16)–(17), is as follows:

$$\llbracket [E, e]\varphi \rrbracket_X = \forall_{R_e^\dagger} \llbracket \varphi \rrbracket_{X \otimes E}, \quad \llbracket \langle E, e \rangle \varphi \rrbracket_X = \exists_{R_e^\dagger} \llbracket \varphi \rrbracket_{X \otimes E}. \quad (24)$$

As in (18), relations  $p_X \circ R_e = p_X \circ q_e \circ i_e^\dagger = i_e \circ i_e^\dagger$  give

$$\forall_{i_e} \circ i_e^{-1} \llbracket \varphi \rrbracket_X = \llbracket \text{Pre}(e) \Rightarrow \varphi \rrbracket_X, \quad \exists_{i_e} \circ i_e^{-1} \llbracket \varphi \rrbracket_X = \llbracket \text{Pre}(e) \wedge \varphi \rrbracket_X, \quad (25)$$

which we may call “static precondition modalities”, as the modal operators of  $i_e \circ i_e^\dagger$ . Then the reduction axioms of DEL can be proven as follows. (The reduction via  $\wedge$  goes since  $\forall_{R_e^\dagger}$  preserves meets, just the same way as in (19); the case of  $\neg$  is similar to (20), albeit more complicated.)

26.  $p_X \circ R_e = i_e \circ i_e^\dagger$  implies the following for atomic  $p$ , by (25) and  $\llbracket p \rrbracket_{X \otimes E} = p_X^{-1} \llbracket p \rrbracket_X$ .

$$\llbracket [E, e]p \rrbracket_X = \forall_{R_e^\dagger} \llbracket p \rrbracket_{X \otimes E} = \forall_{R_e^\dagger} \circ p_X^{-1} \llbracket p \rrbracket_X = \forall_{i_e} \circ i_e^{-1} \llbracket p \rrbracket_X = \llbracket \text{Pre}(e) \Rightarrow p \rrbracket_X.$$

27. For the case of  $\square$ , first note that  $wR_e v p_E e'$  implies  $e = e'$  since  $wR_e v$  implies  $vp_E e$  whereas  $p_E$  is a function. In other words,  $wR_e v p_E e'$  iff  $wR_e v$  and  $e' = e$ . This entails (\*) in the following:

$$\begin{aligned} wR_e; R_{X \otimes E} v &\iff wR_e v' p_X; R_X; p_X^\dagger v \text{ and } wR_e v' p_E; R_E; p_E^\dagger v \text{ for some } v' \in X \otimes E \\ &\stackrel{(*)}{\iff} wR_e; p_X; R_X; p_X^\dagger v \text{ and } eR_E; p_E^\dagger v \\ &\stackrel{(\dagger)}{\iff} wR_e; p_X; R_X; R_e v \text{ for some } e' \in E \text{ such that } eR_E e', \end{aligned}$$

where  $(\dagger)$  holds since  $uR_e v$  iff  $vp_X u$  and  $vp_E e'$ , i.e. iff  $up_X^\dagger v$  and  $e' p_E^\dagger v$ . Thus,

$$R_{X \otimes E} \circ R_e = \left( \bigcup_{eR_E e'} R_{e'} \right) \circ R_X \circ p_X \circ R_e = \left( \bigcup_{eR_E e'} R_{e'} \right) \circ R_X \circ i_e \circ i_e^\dagger.$$

Observe on the other hand that, for a family of relations  $R_i : X \rightarrow Y$  of the same type, we have  $\forall_{(\bigcup_i R_i)^\dagger} = \bigcap_i \circ \forall_{R_i^\dagger}$ . Therefore

$$\begin{aligned} \llbracket [E, e]\square\varphi \rrbracket_X &= \forall_{R_e^\dagger} \circ \forall_{R_{X \otimes E}^\dagger} \llbracket \varphi \rrbracket_{X \otimes E} \\ &= \forall_{i_e} \circ i_e^{-1} \circ \forall_{R_X^\dagger} \circ \bigcap_{eR_E e'} \circ \forall_{R_{e'}^\dagger} \llbracket \varphi \rrbracket_{X \otimes E} = \llbracket \text{Pre}(e) \Rightarrow \square \bigwedge_{eR_E e'} [E, e']\varphi \rrbracket_X. \end{aligned}$$

We conclude this section with a remark on the significance of using the category  $\mathbf{Kr}$ . We reviewed in this section that topological constructions (Subsection 2.4) and their canonical maps play essential rôles in the semantics of PAL and DEL. These constructions take place in  $\mathbf{Kr}$  as opposed to the category  $\mathbf{Kr}_B$ , and the canonical maps are monotone maps of  $\mathbf{Kr}$ , and not bounded morphisms of  $\mathbf{Kr}_B$ . Indeed, for

DEL to show interesting behaviors, the canonical maps—in particular,  $p_X : X \otimes E \rightarrow X$ , which amounts to  $i : \llbracket \sigma \rrbracket_X \hookrightarrow X$  in the case of PAL—must not be bounded morphisms. For, if  $p_X$  is a bounded morphism, then  $\llbracket \varphi \rrbracket_{X \otimes E} = p_X^{-1} \llbracket \varphi \rrbracket_X$  for every  $\varphi$  and not just atomic  $p$  (this entails  $[E, e]\varphi \equiv \text{Pre}(e) \Rightarrow \varphi$  the same way as in (26))—this means that no event can teach agents anything. In other words, for events to teach agents something, they must bring about some change logically, and therefore the maps  $f$  representing them must not have logic-preserving duals  $f^{-1}$ .

## 4 Application: Quantification

This section demonstrates a virtue of our categorical perspective, by showing how to extend DEL to the first order. Our structural approach to DEL and the standard structural approach to first-order logic can be integrated together, simply as two modules, using the methodology of category theory. We will first review how to interpret classical first-order logic in Subsection 4.1, and how to add this first-order structure to Kripke semantics using “Kripke sheaves” in Subsection 4.2. We will then equip this semantics with a DEL-type update in Subsection 4.3, obtaining a new sheaf semantics for first-order DEL.

### 4.1 Classical Semantics in a Slice Category

Here we review how the standard semantics for classical first-order logic goes in the category  $\mathbf{Sets}/X$ , as the non-modal basis of semantics in Subsection 4.2. See [39] for a more general and extensive account.

Let us first recall the definition of *slice category*. Given any category  $\mathbf{C}$ , fix any object  $C$ . Then the slice category  $\mathbf{C}/C$ , “ $\mathbf{C}$  over  $C$ ”, consists of the following:

- Objects are any arrow  $f : D \rightarrow C$  of  $\mathbf{C}$  with the codomain  $C$ .
- Arrows from  $f : D \rightarrow C$  to  $g : E \rightarrow C$  are any arrow  $h : D \rightarrow E$  of  $\mathbf{C}$  such that  $g \circ h = f$ .

In particular, given a set  $X$ ,  $\mathbf{Sets}/X$  is the category of “sets and functions over  $X$ ”:

- Objects, “sets over  $X$ ”, are functions  $\pi : D \rightarrow X$ . For each  $w \in X$  we write  $D_w$  for the inverse image  $\pi^{-1}(\{w\})$ , called the “fiber over  $w$ ”.
- And arrows from  $\pi_1 : D \rightarrow X$  to  $\pi_2 : E \rightarrow X$  are functions  $f : D \rightarrow E$  “over  $X$ ”, meaning that  $\pi_2 \circ f = \pi_1$ , or equivalently that if  $a \in D_w$  then  $f(a) \in E_w$  for the same  $w$ .

We will also later consider a Kripke-structured version of  $\mathbf{Sets}/X$ , viz.  $\mathbf{Kr}/(X, R)$  over a Kripke frame  $(X, R)$ : Its objects and arrows are monotone maps and not just any functions.

Fixing any (nonempty) set  $X$ , the slice category  $\mathbf{Sets}/X$  is used to interpret classical first-order logic as follows. We fix an object  $\pi : D \rightarrow X$  of  $\mathbf{Sets}/X$ , and a surjection  $\pi$  in particular. We then regard  $X$  as a set of worlds and  $D$  as a set of individuals. Each individual  $a \in D$  is assumed to live in a unique world, viz.  $\pi(a) \in X$ . In this sense we may call  $\pi$  a “residence map”. For each world  $w \in X$ , the fiber  $D_w = \pi^{-1}(\{w\})$  is the set of individuals living in  $w$ . In fact, for each  $n \in \mathbb{N}$ , the cartesian product  $D_w^n = D_w \times \cdots \times D_w$  is the set of  $n$ -tuples of individuals living in  $w$ , and the disjoint union of  $D_w^n$  for all  $w \in X$ , i.e. the  $n$ -fold “fibered product” of  $D$  over  $X$ ,

$$D_X^n = \sum_{w \in X} D_w^n = \{(a_1, \dots, a_n) \in D \times \cdots \times D \mid \pi(a_1) = \cdots = \pi(a_n)\},$$

is the set of  $n$ -tuples from the same world, with the projection

$$\pi^n : D_X^n \rightarrow X :: \bar{a} \rightarrow \pi(a_i)$$

mapping an  $n$ -tuple from the same world to that world. (As special cases,  $D_X^1 = D$  and  $D_X^0 = X$ , with  $\pi^1 = \pi : D \rightarrow X$  and  $\pi^0 = 1_X : D \rightarrow D$ .) Categorically speaking, this is to take the  $n$ -fold pullback of  $D$  over  $X$  in **Sets**, or the  $n$ -fold product of  $\pi$  in **Sets**/ $X$ .

One important note regarding the semantics in **Sets**/ $X$  is that it interprets “formulas in contexts”. A context is a (finite) sequence of variables that are all distinct. A formula  $\varphi$  can be in a context  $(x_1, \dots, x_n)$  if no other variables occur freely in  $\varphi$ . It is not assumed that all of  $x_1, \dots, x_n$  actually occur freely in  $\varphi$ ; so, e.g., if  $\varphi$  can be in a context  $(x_1, \dots, x_n)$  then it can also be in  $(x_1, \dots, x_n, x_{n+1}, \dots, x_m)$ . A formula-in-context is a pair of formula and a context it can be in; so, writing  $(x_1, \dots, x_n \mid \varphi)$  presupposes that  $\varphi$  can be in  $(x_1, \dots, x_n)$ . Now, we semantically interpret formulas-in-contexts  $(x_1, \dots, x_n \mid \varphi)$  rather than formulas  $\varphi$ : We regard  $(x_1, \dots, x_n \mid \varphi)$  as an  $n$ -ary predicate that may or may not be true of  $n$ -tuples of individuals  $(a_1, \dots, a_n)$ . Similarly, we interpret terms-in-contexts  $(x_1, \dots, x_n \mid t)$  as mappings of  $n$ -tuples of individuals to individuals. We will write  $\bar{x}$  and  $\bar{a}$  for sequences  $(x_1, \dots, x_n)$  and  $(a_1, \dots, a_n)$ .

In propositional logic, we interpret a sentence  $\sigma$  with  $\llbracket \sigma \rrbracket \subseteq X$ , so that  $w \in \llbracket \sigma \rrbracket$  means that  $\sigma$  is true at  $w$ . Similarly, in the semantics in **Sets**/ $X$ , we interpret a closed sentence  $\sigma$  in the empty context with  $\llbracket \sigma \rrbracket \subseteq X$ . Yet, extending this, we interpret an  $n$ -ary formula-in-context  $(\bar{x} \mid \varphi)$  with  $\llbracket \bar{x} \mid \varphi \rrbracket \subseteq D_X^n$ , so that  $\bar{a} \in \llbracket \bar{x} \mid \varphi \rrbracket$  means that  $\varphi$  is true of individuals  $a_1, \dots, a_n$  in place of  $x_1, \dots, x_n$  (at the world  $\pi(a_i)$ ). The same formula  $\varphi$  in different contexts is true of different tuples: E.g.  $(a, b) \in \llbracket x, y \mid \varphi \rrbracket$  iff  $(b, a) \in \llbracket y, x \mid \varphi \rrbracket$  iff  $(a, b, c) \in \llbracket x, y, z \mid \varphi \rrbracket$  (for any  $c \in D$  such that  $(a, b, c) \in D_X^3$ ).

An interpretation  $\llbracket - \rrbracket$  can be defined inductively, first for terms and then for formulas. In interpreting terms in **Sets**/ $X$ , the core idea is to interpret an  $n$ -ary term-in-context  $(\bar{x} \mid t)$  with an arrow  $\llbracket \bar{x} \mid t \rrbracket : D_X^n \rightarrow D$  in **Sets**/ $X$ , i.e. a function sending  $\bar{a} \in D_w^n$  to  $\llbracket \bar{x} \mid t \rrbracket(\bar{a}) \in D_w$ . To each  $n$ -ary function symbol  $f$ , assign an arrow  $\llbracket f \rrbracket : D_X^n \rightarrow D$  of **Sets**/ $X$ . (This includes  $\llbracket c \rrbracket : X \rightarrow D$  for a constant, i.e. 0-ary function symbol.) Then, for the base case let  $\llbracket \bar{x} \mid f\bar{x} \rrbracket = \llbracket f \rrbracket$ , whereas we also let  $\llbracket \bar{x} \mid x_i \rrbracket = p_i : D_X^n \rightarrow D :: \bar{a} \mapsto a_i$  for each  $i \leq n$ . For inductive steps, define the substitution of terms as follows: Given a term-in-context  $(x_1, \dots, x_n \mid t)$  and terms  $t_1, \dots, t_n$ , we write  $t[t_1, \dots, t_n/x_1, \dots, x_n]$  for the result of substituting  $t_i$  for all the free occurrences of  $x_i$  in  $t$ . Then, given  $\llbracket \bar{x} \mid t \rrbracket$  and  $\llbracket \bar{y} \mid t_i \rrbracket$  for each  $i \leq n$  where  $\bar{y} = (y_1, \dots, y_m)$ , write

$$\llbracket \bar{y} \mid \bar{t} \rrbracket = \langle \llbracket \bar{y} \mid t_1 \rrbracket, \dots, \llbracket \bar{y} \mid t_n \rrbracket \rangle : D_X^m \rightarrow D_X^n :: \bar{b} \mapsto (\llbracket \bar{y} \mid t_1 \rrbracket(\bar{b}), \dots, \llbracket \bar{y} \mid t_n \rrbracket(\bar{b})),$$

and we have

$$\llbracket \bar{y} \mid t[t_1, \dots, t_n/x_1, \dots, x_n] \rrbracket = \llbracket \bar{x} \mid t \rrbracket \circ \llbracket \bar{y} \mid \bar{t} \rrbracket. \quad (28)$$

Now, to each  $n$ -ary relation symbol  $F$ , assign any subset  $\llbracket F \rrbracket \subseteq D_X^n$ , and  $\llbracket \bar{x} \mid F\bar{x} \rrbracket = \llbracket F \rrbracket$ . Inductively,

$$\llbracket \bar{x} \mid \varphi \wedge \psi \rrbracket = \llbracket \bar{x} \mid \varphi \rrbracket \cap \llbracket \bar{x} \mid \psi \rrbracket, \quad \llbracket \bar{x} \mid \neg\varphi \rrbracket = \neg\llbracket \bar{x} \mid \varphi \rrbracket = D_X^n \setminus \llbracket \bar{x} \mid \varphi \rrbracket \quad (29)$$

for Boolean operators. For quantifiers, take a projection  $p : D_X^{n+1} \rightarrow D_X^n :: (\bar{a}, b) \mapsto \bar{a}$  and let

$$\llbracket \bar{x} \mid \forall y. \varphi \rrbracket = \forall_p \llbracket \bar{x}, y \mid \varphi \rrbracket, \quad \llbracket \bar{x} \mid \exists y. \varphi \rrbracket = \exists_p \llbracket \bar{x}, y \mid \varphi \rrbracket; \quad (30)$$

the case of  $n = 0$  is just  $p = \pi : D \rightarrow X$ . Closely connected to quantification is the substitution of terms: Write  $\varphi[\bar{t}/\bar{x}]$  for the result of substituting  $t_i$  for  $x_i$  in  $\varphi$  (this makes sense only if  $t$  is free for  $x$  in  $\varphi$ ). Then the substitution satisfies

$$\llbracket \bar{y} \mid \varphi[t_1, \dots, t_n/x_1, \dots, x_n] \rrbracket = \llbracket \bar{y} \mid \bar{t} \rrbracket^{-1} \llbracket \bar{x} \mid \varphi \rrbracket. \quad (31)$$

As an instance of this, given  $\llbracket \bar{x} \mid \varphi \rrbracket$  we can add a vacuous variable to the context by

$$\llbracket \bar{x}, y \mid \varphi \rrbracket = p^{-1} \llbracket \bar{x} \mid \varphi \rrbracket \quad (32)$$

for the same  $p :: (\bar{a}, b) \mapsto \bar{a}$  as above; and other operations on contexts (e.g. permutation) can be interpreted in similarly obvious ways.

## 4.2 Kripke-Sheaf Semantics

In this subsection we review “Kripke-sheaf semantics” for first-order modal logic. An extensive exposition of this semantics is in [18]. We use the notation and terminology from [26], however, to be consistent with Subsection 4.1.<sup>20</sup>

As to syntax, we take a first-order language—with relation symbols, variables, function symbols and constants—and add  $\Box$  and  $\Diamond$  to it as unary operators that behave just the same way  $\neg$  does. By this we mean in particular that  $\Box(\varphi[t/x])$  (i.e. first substituting  $t$  and then applying  $\Box$ ) and  $(\Box\varphi)[t/x]$  (first applying  $\Box$  and then substituting  $t$ ) are the same formula, just the same way  $\neg(\varphi[t/x])$  and  $(\neg\varphi)[t/x]$  are.

Now, enter

**Definition 1.** A bounded morphism  $\pi : (D, R_D) \rightarrow (X, R_X)$  is called a *Kripke sheaf* over  $(X, R_X)$  if

$$33. aR_D b \pi w \text{ and } aR_D b' \pi w \text{ imply } b = b'. \text{ That is, } (R_D \circ R_D^\dagger) \cap (\pi^\dagger \circ \pi) \subseteq 1_D.$$

We fix one such map and, as we did in Subsection 4.1, regard it as a residence map from the individuals  $D$  to the worlds  $X$ . Then, for each  $n \in \mathbb{N}$ , the set  $D_X^n$  of  $n$ -tuples from the same world comes with the “epistemic” relation  $R_{D_X^n}$  by Fact 3 or by (14) and (15). Categorically, this is to take the  $n$ -fold pullback of  $(D, R_D)$  over  $(X, R_X)$  in  $\mathbf{Kr}$ , or equivalently the  $n$ -fold product of  $\pi$  in the slice category  $\mathbf{Kr}/(X, R_X)$ .

We interpret first-order modal logic with  $\pi$  and other structure in  $\mathbf{Kr}/(X, R_X)$ . The classical base of the logic is interpreted with the underlying, non-Kripke structure in  $\mathbf{Sets}/X$ , just as in Subsection 4.1. The new, modal part is then added to the base using the Kripke structure, as follows: First we require that, for each  $n$ -ary function symbol  $f$ , its interpretation  $\llbracket f \rrbracket : (D_X^n, R_{D_X^n}) \rightarrow (D, R_D)$  be monotone, so that all interpretations  $\llbracket \bar{y} \mid t \rrbracket$  of terms are monotone—i.e., they must be arrows of  $\mathbf{Kr}/(X, R_X)$ . Then we set

$$\llbracket \bar{x} \mid \Box\varphi \rrbracket = \forall_{R_{D_X^n}^\dagger} \llbracket \bar{x} \mid \varphi \rrbracket, \quad \llbracket \bar{x} \mid \Diamond\varphi \rrbracket = \exists_{R_{D_X^n}^\dagger} \llbracket \bar{x} \mid \varphi \rrbracket. \quad (34)$$

In this way, we adopt the following ideas for the semantics.

35. We use a family of Kripke models  $(D_X^n, R_{D_X^n}, \llbracket - \rrbracket)$ , where each  $(D_X^n, R_{D_X^n})$  is the  $n$ -fold product of  $\pi : (D, R_D) \rightarrow (X, R_X)$  in  $\mathbf{Kr}/(X, R_X)$ .
36. Each dual  $(\mathcal{P}(D_X^n), \forall_{R_{D_X^n}^\dagger}, \exists_{R_{D_X^n}^\dagger})$  is a CABA of  $n$ -ary properties governed by (29) and (34).
37. We interpret terms with arrows  $f$  of  $\mathbf{Kr}/(X, R_X)$  between products  $(D_X^n, R_{D_X^n})$ .
38. The CABAOs interact with one another via cross-context operations, which are interpreted, as in (30)–(31), with  $\exists_f \dashv f^{-1} \dashv \forall_f$  of arrows  $f$  of  $\mathbf{Kr}/(X, R_X)$ .

So, let us enter

**Definition 2.** By a *Kripke-sheaf model* we mean a Kripke sheaf  $\pi : (D, R_D) \rightarrow (X, R_X)$  paired with a family of maps  $\llbracket - \rrbracket$  that assigns

- an arrow  $\llbracket f \rrbracket : D_X^n \rightarrow D$  of  $\mathbf{Kr}/(X, R_X)$  to each  $n$ -ary function symbol  $f$ ,
- $\llbracket \bar{y} \mid t \rrbracket : D_X^n \rightarrow D$  to all terms-in-contexts  $(\bar{y} \mid t)$  by  $\llbracket \bar{x} \mid f\bar{x} \rrbracket = \llbracket f \rrbracket$ ,  $\llbracket \bar{x} \mid x_i \rrbracket :: \bar{a} \mapsto a_i$ , and (28),
- any subset  $\llbracket F \rrbracket \subseteq D_X^n$  to each  $n$ -ary relation symbol  $F$ , and
- $\llbracket \bar{x} \mid \varphi \rrbracket \subseteq D_X^n$  to all formulas-in-contexts  $(\bar{x} \mid \varphi)$  by  $\llbracket \bar{x} \mid F\bar{x} \rrbracket = \llbracket F \rrbracket$ , (29)–(32) and (34).

<sup>20</sup>[26] provides a more general semantics using neighborhood structure, but Kripke-sheaf semantics is simply a special case involving Kripke frames; see Section 3 of [26], in particular. It should be noted that the definitions of Kripke sheaf in [18] (Definition 3.6.2) and in [26] (Definition 3.5) only agree for the limited case of reflexive and transitive Kripke frames. Definition 1 in the following is the version in [26].

Definition 2 requires  $\pi$  to be not just a monotone map but moreover a Kripke sheaf, whereas no Kripke sheaves are mentioned in the ideas (35)–(38). The requirement is needed, however, precisely in order for the interaction (38) to behave coherently. Given any  $\llbracket \sigma \rrbracket \subseteq D$ , observe that there are two ways to obtain  $\llbracket y \mid \Box \sigma \rrbracket$  by applying (32) and (34), viz.

$$\llbracket y \mid \Box \sigma \rrbracket = \forall_{R_D^\dagger} \llbracket y \mid \sigma \rrbracket = \forall_{R_D^\dagger} \circ \pi^{-1} \llbracket \sigma \rrbracket, \quad \llbracket y \mid \Box \sigma \rrbracket = \pi^{-1} \llbracket \Box \sigma \rrbracket = \pi^{-1} \circ \forall_{R_X^\dagger} \llbracket \sigma \rrbracket.$$

So the well-definedness of  $\llbracket - \rrbracket$ , along with (32) and (34), requires that  $\forall_{R_D^\dagger} \circ \pi^{-1} = \pi^{-1} \circ \forall_{R_X^\dagger}$ , or dually  $\pi \circ R_D = R_X \circ \pi$ , i.e. that  $\pi$  be a bounded morphism. Indeed, any map  $\llbracket \bar{y} \mid \bar{t} \rrbracket$  involved in (31) must be a bounded morphism. Recall that our syntax has  $\Box(\varphi[\bar{t}/\bar{x}]) = (\Box\varphi)[\bar{t}/\bar{x}]$ . This means that, for  $\llbracket - \rrbracket$  to be well-defined, we need  $\llbracket \bar{y} \mid \Box(\varphi[\bar{t}/\bar{x}]) \rrbracket = \llbracket \bar{y} \mid (\Box\varphi)[\bar{t}/\bar{x}] \rrbracket$ , both sides giving the same interpretation to the same formula  $\Box\varphi[\bar{t}/\bar{x}]$ . So, given  $\llbracket \bar{x} \mid \varphi \rrbracket \subseteq D_X^n$  and  $\llbracket \bar{t} \rrbracket = \llbracket \bar{y} \mid \bar{t} \rrbracket : D_X^m \rightarrow D_X^n$ , (31) and (34) imply

$$\forall_{R_{D_X^m}^\dagger} \circ \llbracket \bar{t} \rrbracket^{-1} \llbracket \bar{x} \mid \varphi \rrbracket = \llbracket \bar{y} \mid \Box(\varphi[\bar{t}/\bar{x}]) \rrbracket = \llbracket \bar{y} \mid (\Box\varphi)[\bar{t}/\bar{x}] \rrbracket = \llbracket \bar{t} \rrbracket^{-1} \circ \forall_{R_{D_X^n}^\dagger} \llbracket \bar{x} \mid \varphi \rrbracket.$$

Thus, the well-definedness of  $\llbracket - \rrbracket$ , along with (31) and (34), again requires that  $\llbracket \bar{t} \rrbracket$  be a bounded morphism.<sup>21</sup> Yet, all maps involved in Definition 2 are indeed guaranteed to be bounded morphisms, by

**Fact 4.** If  $\pi : D \rightarrow X$  is a Kripke sheaf, then so is every  $\pi^n : D_X^n \rightarrow X$ . Moreover, given two Kripke sheaves  $\pi_D : D \rightarrow X$  and  $\pi_E : E \rightarrow X$ , any monotone map  $f : D \rightarrow E$  over  $X$  (i.e. satisfying  $\pi_E \circ f = \pi_D$ ) is also a Kripke sheaf (and hence a bounded morphism). On the other hand,  $\pi$  is a Kripke sheaf iff both  $\pi$  and the “diagonal map”  $\Delta : D \rightarrow D_X^2 :: a \mapsto (a, a)$  are bounded morphisms.<sup>22</sup>

In short, the simple combination of (28)–(32), for classical first-order logic, and (29) and (34), for propositional modal logic, is made possible by Kripke sheaves and Fact 4. And this simple combination makes the logic of Kripke-sheaf semantics the simple union of classical first-order logic and modal logic.

**Fact 5.** Let **FOK** be the first-order modal logic that consists of all the rules and axioms of classical first-order logic, and the rules and axioms of propositional modal logic **K**. Then **FOK** is sound and complete with respect to the Kripke-sheaf models. The same holds with **S4** (or **S5**, respectively) in place of **K**, with respect to the subclass of Kripke-sheaf models over preorders (or equivalence relations).<sup>23</sup>

### 4.3 First-Order Dynamic Epistemic Logic

In Subsection 4.2 we saw how the Kripke-sheaf structure extended the modal logic of a Kripke model to the first order. We will now lay out how the same structure can extend the product update of Kripke models to the first order.<sup>24</sup> One remark is in order: We saw in Sections 2 and 3 that, whereas bounded morphisms play a more prominent rôle than merely monotone maps in the semantics of static modal logic, merely monotone maps are essential in the semantics of DEL. This theme recurs in this subsection.

<sup>21</sup>From a perspective of categorical logic, one often takes (31), for all  $\varphi$ , as part of the definition of a model, rather than a derived fact about the model. It is from this perspective that we describe the situation as a matter of well-definedness of the model. One could also see the same situation as a matter of deriving (31) from its atomic case using a property of bounded morphisms; we acknowledge an anonymous reviewer for this perspective. One could of course choose to reject (31) or (34), or even to use a syntax without  $\Box(\varphi[t/x]) = (\Box\varphi)[t/x]$ . (These options, needless to say, would make Fact 5 unavailable to one’s semantics.) A notable case of rejecting (34) is the counterpart theory in [34], which restricts (34) to the case where all the variables in  $\bar{x}$  actually occur freely in  $\varphi$ .

<sup>22</sup>See Facts 4.2, 4.4, and 4.6 in [26].

<sup>23</sup>See, e.g., Corollary 6.1.24 of [18].

<sup>24</sup>A sheaf semantics for first-order PAL was given (in a more general, neighborhood setting) in [27]. A first-order extension of PAL was also given briefly in [35], which, however, used constant domains to interpret quantification. See footnote 2 as well.

In Subsection 4.2, we reviewed the fact that static first-order modal logic needed Kripke sheaves to make sure all the maps involved were bounded morphisms. In our new semantics for first-order DEL, however, the structure of the category  $\mathbf{Kr}$  of monotone maps will play a central rôle again.

Let  $(\pi : (D, R_D) \rightarrow (X, R_X), \llbracket - \rrbracket_\pi)$  be a Kripke-sheaf model, and  $(E, R_E)$  be a Kripke frame. We regard the latter as an event model, and assume that preconditions  $\text{Pre}(e)$  for  $e \in E$  are all (closed) sentences, so that  $\llbracket \text{Pre}(e) \rrbracket_\pi$  makes sense and  $\llbracket \text{Pre}(e) \rrbracket_\pi \subseteq X$ . Then  $(X, R_X)$  is product-updated with  $(E, R_E)$  into  $(X \otimes E, R_{X \otimes E})$ . For the first-order structure, we moreover “pullback-update”  $(\pi, \llbracket - \rrbracket_\pi)$ , by pulling everything back along the projection  $p_X : X \otimes E \rightarrow X$ . Recall that  $(\pi, \llbracket - \rrbracket_\pi)$  uses the structure of the slice category  $\mathbf{Kr}/(X, R_X)$ ; hence  $p_X$  induces a pullback functor  $p_X^* : \mathbf{Kr}/(X, R_X) \rightarrow \mathbf{Kr}/(X \otimes E, R_{X \otimes E})$ . So we apply this to obtain an updated residence map  $\pi_{X \otimes E} = p_X^* \pi : D_{X \otimes E} \rightarrow X \otimes E$ , and to obtain  $\llbracket - \rrbracket_{\pi_{X \otimes E}}$  from  $\llbracket f \rrbracket_{\pi_{X \otimes E}} = p_X^* \llbracket f \rrbracket_\pi$  for function symbols  $f$  and  $\llbracket F \rrbracket_{\pi_{X \otimes E}} = p_X^* \llbracket F \rrbracket_\pi$  for relation symbols  $F$ . We need to note that the structure of  $\mathbf{Kr}$  is essential for the pullback update. Pullbacks are taken in the category  $\mathbf{Kr}$  of monotone maps in general as opposed to bounded morphisms, and along the map  $p_X : X \otimes E \rightarrow X$ , which, as seen in Subsection 3.2, must not be a bounded morphism for DEL to show interesting behaviors.

Here is an explicit description of the pullback update:

- Using the notation  $D_w = \pi^{-1}(\{w\})$ , the pullback of  $\pi^n : D_X^n \rightarrow X :: \bar{a} \mapsto \pi(a_i)$  along  $p_X : X \otimes E \rightarrow X :: (w, e) \mapsto w$  has the domain

$$\sum_{(w,e) \in X \otimes E} D_w^n = \{(\bar{a}, e) \in D_X^n \times E \mid \pi^n(\bar{a}) \in \llbracket \text{Pre}(e) \rrbracket_\pi\},$$

for which we write  $D_{X \otimes E}^n$ , and projections

$$\begin{aligned} \pi_{X \otimes E}^n &= p_X^* \pi^n : D_{X \otimes E}^n \rightarrow X \otimes E :: (\bar{a}, e) \mapsto (\pi^n(\bar{a}), e), \\ p_{D_X^n} &: D_{X \otimes E}^n \rightarrow D_X^n :: (\bar{a}, e) \mapsto \bar{a}. \end{aligned}$$

It also comes with another projection  $p_{E,n} : D_{X \otimes E}^n \rightarrow E :: (\bar{a}, e) \mapsto e$ . The “epistemic” relation  $R_{D_{X \otimes E}^n}$  on  $D_{X \otimes E}^n$  is an initial lift, viz.

$$\begin{aligned} R_{D_{X \otimes E}^n} &= (p_{D_X^n}^\dagger \circ R_{D_X^n} \circ p_{D_X^n}) \cap (p_{E,n}^\dagger \circ R_E \circ p_{E,n}), \\ \text{i.e. } (\bar{a}, e_1) R_{D_{X \otimes E}^n} (\bar{b}, e_2) &\iff a_1 R_D b_1, \dots, a_n R_D b_n \text{ and } e_1 R_E e_2. \end{aligned}$$

- For an  $n$ -ary function symbol  $f$ , we have  $\llbracket f \rrbracket_\pi : D_X^n \rightarrow D$  and then

$$\llbracket f \rrbracket_{\pi_{X \otimes E}} = p_X^* \llbracket f \rrbracket_\pi : D_{X \otimes E}^n \rightarrow D_{X \otimes E} :: (\bar{a}, e) \mapsto (\llbracket f \rrbracket_\pi(\bar{a}), e).$$

- For an  $n$ -ary relation symbol  $F$ , we have  $\llbracket F \rrbracket_\pi \subseteq D_X^n$  and then

$$\llbracket F \rrbracket_{\pi_{X \otimes E}} = p_X^* \llbracket F \rrbracket_\pi = p_{D_X^n}^{-1} \llbracket F \rrbracket_\pi = \{(\bar{a}, e) \in D_{X \otimes E}^n \mid \bar{a} \in \llbracket F \rrbracket_\pi\} \subseteq D_{X \otimes E}^n.$$

The pullback update indeed updates a Kripke-sheaf model to another:

**Theorem 2.** *Given a Kripke-sheaf model  $(\pi, \llbracket - \rrbracket_\pi)$ , its pullback update  $(\pi_{X \otimes E}, \llbracket - \rrbracket_{\pi_{X \otimes E}})$  along  $p_X : X \otimes E \rightarrow X$  is a Kripke-sheaf model.*

*Proof.* As in Fact 4, both  $\pi$  and the diagonal map  $\Delta$  of  $\pi$  are bounded morphisms, and hence Theorem 1 implies that both  $\pi_{X \otimes E} = p_X^* \pi$  and  $p_X^* \Delta$  are bounded morphisms. Yet  $p_X^* \Delta$  is the diagonal map of  $\pi_{X \otimes E}$ , since the pullback functor  $p_X^*$  preserves finite limits. Therefore  $\pi_{X \otimes E}$  is a Kripke sheaf by Fact 4. Moreover, for each  $n \in \mathbb{N}$ ,  $p_X^* \pi^n$  is the  $n$ -fold product of  $\pi_{X \otimes E}$  over  $X \otimes E$ , since  $p_X^*$  preserves finite limits.  $\square$

Now we have two Kripke-sheaf models,  $(\pi, \llbracket - \rrbracket_\pi)$  before update and  $(\pi_{X \otimes E}, \llbracket - \rrbracket_{\pi_{X \otimes E}})$  after, and we can use relations between them to interpret the DEL operators  $[E, e]$  and  $\langle E, e \rangle$ . Here is a key idea: As in (35)–(36), each sheaf model has a Kripke model for  $n$ -ary properties,  $D_X^n$  and  $D_{X \otimes E}^n$ ; so we treat  $D_X^n$  and  $D_{X \otimes E}^n$  as the product-update structure of Subsection 3.2 that interprets the application of  $[E, e]$  and  $\langle E, e \rangle$  to  $n$ -ary formulas-in-contexts. Since  $\llbracket \bar{x} \mid \text{Pre}(e) \rrbracket_\pi = (\pi^n)^{-1} \llbracket \text{Pre}(e) \rrbracket$ , observe

$$D_{X \otimes E}^n = \sum_{e \in E} \llbracket \bar{x} \mid \text{Pre}(e) \rrbracket_\pi = \{ (\bar{a}, e) \in D_X^n \times E \mid \bar{a} \in \llbracket \bar{x} \mid \text{Pre}(e) \rrbracket_\pi \}$$

and note the similarity to (22). We moreover have canonical maps as with (22), viz. the projection  $p_{D_X^n} : D_{X \otimes E}^n \rightarrow D_X^n$  above and, for each  $e \in E$ ,

- The inclusion map  $i_e^n : \llbracket \bar{x} \mid \text{Pre}(e) \rrbracket_\pi \hookrightarrow D_X^n$ .
- The coproduct injection  $q_e^n : \llbracket \bar{x} \mid \text{Pre}(e) \rrbracket_\pi \rightarrow D_{X \otimes E}^n :: \bar{a} \mapsto (\bar{a}, e)$ .

These maps tabulate a relation,  $R_e^n = q_e^n \circ i_e^{n\dagger} : D_X^n \rightarrow D_{X \otimes E}^n$ , which is dual to the two maps

$$\forall_{R_e^{n\dagger}} = \forall_{i_e^n} \circ (q_e^n)^{-1}, \exists_{R_e^{n\dagger}} = \exists_{i_e^n} \circ (q_e^n)^{-1} : \mathcal{P}(D_{X \otimes E}^n) \rightarrow \mathcal{P}(D_X^n).$$

These then interpret  $[E, e]$  and  $\langle E, e \rangle$  applied to  $n$ -ary formulas-in-contexts  $(\bar{x} \mid \varphi)$ , i.e.,

$$\llbracket \bar{x} \mid [E, e]\varphi \rrbracket_\pi = \forall_{R_e^{n\dagger}} \llbracket \bar{x} \mid \varphi \rrbracket_{\pi_{X \otimes E}}, \quad \llbracket \bar{x} \mid \langle E, e \rangle \varphi \rrbracket_\pi = \exists_{R_e^{n\dagger}} \llbracket \bar{x} \mid \varphi \rrbracket_{\pi_{X \otimes E}},$$

which is just an “in context” version of (24).

This defines our sheaf semantics for first-order DEL—but we need to check its well-definedness, similarly to the remark following Definition 2. That is, we need

$$\begin{aligned} \forall_{R_e^{m\dagger}} \circ \llbracket \bar{t} \rrbracket_{\pi_{X \otimes E}}^{-1} \llbracket \bar{x} \mid \varphi \rrbracket_{\pi_{X \otimes E}} &= \llbracket \bar{y} \mid [E, e](\varphi[\bar{t}/\bar{x}]) \rrbracket_\pi \\ &= \llbracket \bar{y} \mid ([E, e]\varphi)[\bar{t}/\bar{x}] \rrbracket_\pi = \llbracket \bar{t} \rrbracket_\pi^{-1} \circ \forall_{R_e^{m\dagger}} \llbracket \bar{x} \mid \varphi \rrbracket_{\pi_{X \otimes E}}, \end{aligned}$$

and similarly for  $\langle E, e \rangle$ . Yet these are the case because  $\llbracket \bar{t} \rrbracket_{\pi_{X \otimes E}} \circ R_e^m = R_e^m \circ \llbracket \bar{t} \rrbracket_\pi$  by

**Theorem 3.** For any arrow  $f : D_X^m \rightarrow D_X^n$  of  $\mathbf{Kr}/(X, R_X)$ ,

$$p_X^* f \circ R_e^m = R_e^n \circ f, \quad (p_X^* f)^\dagger \circ R_e^n = R_e^m \circ f^\dagger.$$

*Proof.* This follows from Corollary 3 since the following squares are both pullbacks in **Sets**.

$$\begin{array}{ccccc} D_{X \otimes E}^m & \xleftarrow{q_e^m} & (\pi^m)^{-1} \llbracket \text{Pre}(e) \rrbracket_\pi & \xleftarrow{i_e^m} & D_X^m \\ p_X^* f \downarrow & & \lrcorner \downarrow \lrcorner & & \downarrow f \\ D_{X \otimes E}^n & \xleftarrow{q_e^n} & (\pi^n)^{-1} \llbracket \text{Pre}(e) \rrbracket_\pi & \xleftarrow{i_e^n} & D_X^n \end{array} \quad \square$$

Now, the semantics validates all the reduction axioms of propositional DEL, simply because  $D_{X \otimes E}^n$  is just the product update of  $D_X^n$  with  $(E, R_E)$ . One more reduction axiom is needed, however—viz. for quantifiers. And here it is:

$$[E, e]\forall y. \varphi \equiv \forall y. [E, e]\varphi. \quad (39)$$



*Proof.* We show the validity of (39). Let  $p : D_X^{n+1} \rightarrow D_X^n :: (\bar{a}, b) \mapsto \bar{a}$ . Then Theorem 3 implies  $(p_X^* p)^\dagger \circ R_e^n = R_e^{n+1} \circ p^\dagger$ , and dually  $\forall_{R_e^{n+1}} \circ \forall_{p_X^* p} = \forall_p \circ \forall_{R_e^n}$ . Therefore

$$\llbracket \bar{x} \mid [E, e] \forall y. \varphi \rrbracket_\pi = \forall_{R_e^{n+1}} \circ \forall_{p_X^* p} \llbracket \bar{x}, y \mid \varphi \rrbracket_{\pi_{X \otimes E}} = \forall_p \circ \forall_{R_e^n} \llbracket \bar{x}, y \mid \varphi \rrbracket_{\pi_{X \otimes E}} = \llbracket \bar{x} \mid \forall y. [E, e] \varphi \rrbracket_\pi. \quad \square$$

This now gives a completeness result extending Fact 5 by the standard method of reduction.

**Theorem 4.** *Let FODEL-K be the first-order modal logic that consists of FOK, all the reduction axioms of propositional DEL, and (39). Then FODEL-K is sound and complete with respect to the Kripke-sheaf models with pullback updates. The versions with S4 and S5 in place of K hold with respect to the obvious subclasses of Kripke-sheaf models.*

## 5 Connections to Preceding Approaches

There have been approaches to modal logic and DEL that take advantage of concepts and methods of category theory in different ways from our approach. This section discusses connections between some of these approaches and ours.<sup>25</sup>

Semantics of modal logic S4 shows various categorical structures. A Kripke frame  $(X, \lesssim)$  for S4 is a preorder, and hence itself a category. Also, the family  $\mathcal{O}X$  of  $\lesssim$ -upward closed subsets of  $X$  forms a topology on  $X$ , and hence a category. Moreover, the interior operation  $\text{int} : \mathcal{P}X \rightarrow \mathcal{O}X$  of this topology is right adjoint to the inclusion  $i : \mathcal{O}X \hookrightarrow \mathcal{P}X$ , so that  $\square = i \circ \text{int}$  is the comonad of the adjunction.<sup>26</sup> The notion of (Kripke) sheaf lifts all these structures to the first order: A Kripke sheaf over a preorder  $(X, \lesssim)$  is equivalently a “presheaf” on the category  $(X, \lesssim)$ , an “étale space” over the space  $(X, \mathcal{O}X)$ , and a “sheaf” on the category  $\mathcal{O}X$ .<sup>27</sup> Moreover, the adjunction  $i \dashv \text{int}$  is lifted to a “geometric morphism”  $i^* \dashv i_*$  from **Sets**/ $X$  to the “topos” of sheaves over  $\mathcal{O}X$ , so that its comonad  $i^* \circ i_*$  induces  $\square : \mathcal{P}D \rightarrow \mathcal{P}D$  for every Kripke sheaf  $\pi : (D, \lesssim_D) \rightarrow (X, \lesssim)$ .<sup>28</sup> Not all these categorical structures carry over to the general (i.e. non-S4) Kripke semantics. It will be interesting, however, to investigate how to integrate them with DEL updates, given that epistemic relations are normally assumed to be preorders. In fact, given a monotone map  $f : (X, \lesssim_X) \rightarrow (Y, \lesssim_Y)$  of preorders, the pullback functor  $f^*$  (which plays a key rôle in the pullback update of Subsection 4.3) has a right adjoint  $f_*$ , and  $f^* \dashv f_*$  is a typical example of geometric morphism, from the topos of Kripke sheaves over  $(X, \lesssim_X)$  to those over  $(Y, \lesssim_Y)$ .

A categorical approach that covers the entire Kripke semantics (for static modal logic) is given by coalgebras (see, e.g., [38, 13, 30, 23]). The category **Rel** of relations is the “Kleisli category” of the “powerset monad”  $\mathcal{P} : \mathbf{Sets} \rightarrow \mathbf{Sets}$ , meaning, among other things, that the relations  $R : X \twoheadrightarrow Y$  correspond 1–1 to the functions  $r : X \rightarrow \mathcal{P}Y$ .<sup>29</sup> Indeed, the powerset monad is precisely the duality  $\exists_- : \mathbf{Rel} \rightarrow \mathbf{CABA}_\vee$  restricted to **Sets** (and followed by the forgetful  $U : \mathbf{CABA}_\vee \rightarrow \mathbf{Sets}$ ). The correspondence implies that the Kripke frames  $(X, R : X \twoheadrightarrow X)$  are exactly the coalgebras  $r : X \rightarrow \mathcal{P}X$  for the endofunctor  $\mathcal{P}$ . Their homomorphisms, from  $r_X : X \rightarrow \mathcal{P}X$  to  $r_Y : Y \rightarrow \mathcal{P}Y$ , are normally defined as functions  $f : X \rightarrow Y$  satisfying  $\exists_f \circ r_X = r_Y \circ f$ , which amounts to (10),  $f \circ R_X = R_Y \circ f$ , for the corresponding relations  $R_X$  and

<sup>25</sup>We thank anonymous reviewers for references, and for their suggestions that the connections should be discussed.

<sup>26</sup>See Section 10.4 of [3] and Subsection 5.1.1 of [23] for comonads. In fact, instead of a poset  $\mathcal{P}X$  one can take a general category **C** and a comonad  $\square$  on **C** to interpret S4 (perhaps with a non-modal base weaker than classical); see e.g. [2].

<sup>27</sup>See Chapters I through III of [37] for these concepts.

<sup>28</sup>See Chapter VII of [37] for geometric morphisms in general, and Section 5.2 of [4] for the geometric-morphism interpretation of FOS4.

<sup>29</sup>This correspondence can also be described as between  $R : X \times Y \rightarrow \mathbf{2}$  and  $r : X \rightarrow (Y \rightarrow \mathbf{2})$ . See Chapter VI of [36], Chapter 10 of [3], and Chapter 5 of [23] for monads and their Kleisli categories, and  $\mathcal{P}$  and **Rel** as an example. Fact 1 can then be read as stating that  $\exists_-$  is a “comparison functor” that presents **Rel** as the category  $\mathbf{CABA}_\vee$  of free algebras of  $\mathcal{P}$ .

$R_Y$ . Therefore, in the coalgebraic approach to Kripke semantics,  $\mathbf{Coalg}(\mathcal{P})$ , the category of coalgebras and their homomorphisms normally considered, is—like the category  $\mathbf{CABAO}$  of CABAOs and their homomorphisms—equivalent to the category  $\mathbf{Krb}$  of bounded morphisms. In this article, on the other hand, we emphasized the significance of the topological category  $\mathbf{Kr}$  of monotone maps for DEL.<sup>30</sup>

There have in fact been algebraic [31] and coalgebraic [5, 14] approaches to DEL. In particular, the algebraic approach by Kurz and Palmigiano [31] uses ideas closely related to those in Section 3 of this article: They observe that the product update  $X \otimes E$  is a subframe of the coproduct  $X \times E = \sum_{e \in E} X$ , and study the dual structure, i.e. a quotient of the product  $\prod_{e \in E} \mathcal{P}(X)$ .<sup>31</sup> Kurz and Palmigiano are well aware that these constructions do not take place in  $\mathbf{Krb}$  or  $\mathbf{CABAO}$  but rather in  $\mathbf{Kr}$  and  $\mathbf{CABAO}_C$ . They stop short, however, of studying  $\mathbf{Kr}$  or  $\mathbf{CABAO}_C$ , saying that “for these dual characterizations to be defined, an *a priori* specification of the fully fledged category-theoretic environment in which these constructions are taken is actually not needed” ([31], 2). We, in contrast, work under the philosophy that, when one finds a good heuristic that leads to a new result, they should study the heuristic itself and shape it into a theory that yields more results systematically. The point of Section 4 was to demonstrate how to put to use more structures in  $\mathbf{Kr}$ . It should also be stressed that we use one more category, viz.  $\mathbf{Rel}$ , and take essential advantage of the fundamental relation-modality dualities of Subsection 2.2, and not just the derivative dualities of Subsection 2.3 between Kripke frames and CABAOs.

## 6 Conclusion and Future Work

In this paper we have recast the standard semantics of dynamic epistemic logic (DEL) in categorical terms and shed new structural light on it. It should be clear by now how conceptually powerful the new way of applying categorical method is: As demonstrated by our new semantics for first-order DEL, our categorical, structural perspective tightly connects what we want (or need) logically or syntactically and what we need (or want) semantically.

Our new application of the categorical methodology promises to be helpful on multiple fronts of the study of DEL. Naturally expected future work is to extend our approach to more vocabulary (e.g. common knowledge or  $\mu$ -calculus), more types of logic (e.g. higher-order DEL or typed DEL), more structures (e.g. probability), and more general settings (e.g. intuitionistic or constructive modal logic). Various updates can be expressed as functors between categories of models, and these expressions are expected to help characterize properties of updates such as the preservation of constructions or the admitting of reduction axioms. As mentioned in Section 5, the case of  $\mathbf{S4}$  can be formulated in terms of toposes. Or our structural, topological ideas on the category  $\mathbf{Kr}$  of monotone maps for DEL can be used

---

<sup>30</sup>One can of course express  $\mathbf{Kr}$  with coalgebras, by defining a weaker notion of homomorphism, corresponding to monotone maps—i.e., a function  $f : X \rightarrow Y$  is “continuous” from  $r_X : X \rightarrow \mathcal{P}X$  to  $r_Y : Y \rightarrow \mathcal{P}Y$  if  $\exists_f \circ r_X \leq r_Y \circ f$  (i.e.  $\exists_f \circ r_X(x) \subseteq r_Y \circ f(x)$  for all  $x \in X$ ). On the other hand, Kripke sheaves can be defined within  $\mathbf{Coalg}(\mathcal{P})$ . One can rewrite (33) as a homomorphism  $\pi : D \rightarrow X$  from  $r_D : D \rightarrow \mathcal{P}D$  to  $r_X : X \rightarrow \mathcal{P}X$  satisfying

- for each  $a \in D$ , the restriction of  $\pi$  to  $r_D(a)$  is an injection.

Or it may be better to use the characterization in Fact 4—i.e.,  $\pi$  is a Kripke sheaf iff both  $\pi$  and  $\Delta$  are homomorphisms. See Fact 4.2 of [26]. The latter definition can indeed be extended to more kinds of coalgebras and not just Kripke frames.

<sup>31</sup>It is therefore the maps  $i$  and  $q'_e$  in (23) that play a central rôle in [31]. In contrast, we put more emphasis on  $q_e$  and  $i_e$ , though  $R_e = q_e \circ i_e^\dagger = i^\dagger \circ q'_e$  as noted on p. 362. Also, in our treatment, the characterization of  $X \times E$  as a product plays a key role as well, since the Kripke frame on  $X \times E$  is the product of  $X$  and  $E$ , but not the coproduct of  $(X)_{e \in E}$ , in  $\mathbf{Kr}$ . Moreover, we treat  $\text{Pre}(e) \Rightarrow -$  and  $\text{Pre}(e) \wedge -$  as the modal operators of  $i_e \circ i_e^\dagger$ , a perspective that then enables us to prove the reduction axioms (21) and (27) directly by the relation-modality duality  $\forall_{-\dagger}$ . This should be contrasted to the treatment of  $\text{Pre}(e) \Rightarrow -$  and  $\text{Pre}(e) \wedge -$  in proofs in Section 7 of [31].

to augment the coalgebraic generalization of the subcategory  $\mathbf{Krb}$  of bounded morphisms. One may also find, e.g., (39) too strong for their purpose, and hence need to replace the pullback update with a more flexible idea. Furthermore, although we formulated a categorical semantics, we did not mention a crucial aspect of categorical logic—viz. an interpretation  $\llbracket - \rrbracket$  as a homomorphism. To cover this aspect we need to define a “syntactic category” for DEL; this will then lead to a new theory of duality.

## References

- [1] Jiří Adámek, Horst Herrlich & George E. Strecker (1990): *Abstract and Concrete Categories: The Joy of Cats*. John Wiley and Sons.
- [2] Natasha Alechina, Michael Mendler, Valeria de Paiva & Eike Ritter (2001): *Categorical and Kripke Semantics for Constructive S4 Modal Logic*. In Laurent Fribourg, editor: *Computer Science Logic (CSL'01)*, LNCS 2142, Springer, pp. 292–307, doi:10.1007/3-540-44802-0\_21.
- [3] Steve Awodey (2010): *Category Theory*, second edition. Oxford University Press.
- [4] Steve Awodey & Kohei Kishida (2008): *Topology and Modality: The Topological Interpretation of First-Order Modal Logic*. *Review of Symbolic Logic* 1, pp. 146–166, doi:10.1017/S1755020308080143.
- [5] Alexandru Baltag (2003): *A Coalgebraic Semantics for Epistemic Programs*. *Electronic Notes in Theoretical Computer Science* 82, pp. 17–37, doi:10.1016/S1571-0661(04)80630-3.
- [6] Alexandru Baltag & Lawrence Moss (2004): *Logics for Epistemic Programs*. *Synthese* 139, pp. 165–224, doi:10.1023/B:SYNT.0000024912.56773.5e.
- [7] Alexandru Baltag, Lawrence Moss & Sławomir Solecki (1998): *The Logic of Public Announcements, Common Knowledge and Private Suspicions*. In Itzhak Gilboa, editor: *Proceedings of the 7th Conference on Theoretical Aspects of Rationality and Knowledge (TARK VII)*, pp. 43–56. Available at [http://www.tark.org/proceedings/tark\\_jul22\\_98/p43-baltag.pdf](http://www.tark.org/proceedings/tark_jul22_98/p43-baltag.pdf).
- [8] Johan van Benthem (2011): *Logical Dynamics of Information and Interaction*. Cambridge University Press, doi:10.1017/CBO9780511974533.
- [9] Johan van Benthem, Jan van Eijck & Barteld Kooi (2006): *Logics of Communication and Change*. *Information and Computation* 204, pp. 1620–1662, doi:10.1016/j.ic.2006.04.006.
- [10] Patrick Blackburn, Maarten de Rijke & Yde Venema (2001): *Modal Logic*. Cambridge University Press, doi:10.1017/CBO9781107050884.
- [11] Aurelio Carboni & Robert F. C. Walters (1987): *Cartesian Bicategories I*. *Journal of Pure and Applied Algebra* 49, pp. 11–32, doi:10.1016/0022-4049(87)90121-6.
- [12] Alexander Chagrov & Michael Zakharyashev (1997): *Modal Logic*. Clarendon Press.
- [13] Corina Cîrstea, Alexander Kurz, Dirk Pattinson, Lutz Schröder & Yde Venema (2011): *Modal Logics are Coalgebraic*. *The Computer Journal* 54, pp. 31–41, doi:10.1093/comjnl/bxp004.
- [14] Corina Cîrstea & Mehrnoosh Sadrzadeh (2007): *Coalgebraic Epistemic Update Without Change of Model*. In Till Mossakowski, Ugo Montanari & Magne Haveraaen, editors: *Algebra and Coalgebra in Computer Science (CALCO 2007)*, LNCS 4642, Springer, pp. 158–172, doi:10.1007/978-3-540-73859-6\_11.
- [15] Bob Coecke & Éric Oliver Paquette (2011): *Categories for the Practising Physicist*. In Bob Coecke, editor: *New Structures for Physics*, Springer, pp. 173–286, doi:10.1007/978-3-642-12821-9\_3.
- [16] Hans van Ditmarsch, Wiebe van der Hoek & Barteld Kooi (2008): *Dynamic-Epistemic Logic*. Springer, doi:10.1007/978-1-4020-5839-4.
- [17] Peter J. Freyd & Andre Scedrov (1990): *Categories, Allegories*. North-Holland.
- [18] Dov M. Gabbay, Valentin Shehtman & Dimitrij Skvortsov (2009): *Quantification in Nonclassical Logic*. 1, Elsevier.

- [19] Silvio Ghilardi (2010): *Continuity, Freeness, and Filtrations*. *Journal of Applied Non-Classical Logics* 20, pp. 193–217, doi:10.3166/jancl.20.193-217.
- [20] David Harel (1979): *First-Order Dynamic Logic*. Springer, doi:10.1007/3-540-09237-4.
- [21] David Harel, Dexter Kozen & Jerzy Tiuryn (2000): *Dynamic Logic*. MIT Press.
- [22] Claudio Hermida (2011): *A Categorical Outlook on Relational Modalities and Simulations*. *Information and Computation* 209, pp. 1505–1517, doi:10.1016/j.ic.2010.09.009.
- [23] Bart Jacobs (2016): *Introduction to Coalgebra: Towards Mathematics of States and Observation*. Cambridge University Press, doi:10.1017/CBO9781316823187.
- [24] André Joyal & Myles Tierney (1984): *An Extension of the Galois Theory of Grothendieck*. American Mathematical Society.
- [25] Burghard von Karger (1998): *Temporal Algebra*. *Mathematical Structures in Computer Science* 8, pp. 277–320, doi:10.1017/S0960129598002540.
- [26] Kohei Kishida (2011): *Neighborhood-Sheaf Semantics for First-Order Modal Logic*. *Electronic Notes in Theoretical Computer Science* 278, pp. 129–143, doi:10.1016/j.entcs.2011.10.011.
- [27] Kohei Kishida (2013): *Public Announcements under Sheaves*. In Yoichi Motomura, Alastair Butler & Daisuke Bekki, editors: *New Frontiers in Artificial Intelligence (JSAI-isAI 2012 Workshops)*, LNCS 7856, Springer, pp. 96–108, doi:10.1007/978-3-642-39931-2\_8.
- [28] Kohei Kishida (forthcoming): *Categories and Modalities*. In Elaine M. Landry, editor: *Categories for the Working Philosopher*, Oxford University Press.
- [29] Barteld Kooi (2007): *Dynamic Term-Modal Logic*. In: *A Meeting of the Minds: Proceedings of the Workshop on Logic, Rationality and Interaction, Beijing, 2007*, College Publications, pp. 173–185.
- [30] Clemens Kupke & Dirk Pattinson (2011): *Coalgebraic Semantics of Modal Logics: An Overview*. *Theoretical Computer Science* 412, pp. 5070–5094, doi:10.1016/j.tcs.2011.04.023.
- [31] Alexander Kurz & Alessandra Palmigiano (2013): *Epistemic Updates on Algebras*. *Logical Methods in Computer Science* 9(4):17, doi:10.2168/LMCS-9(4:17)2013.
- [32] Stephen Lack (2010): *A 2-Categories Companion*. In John C. Baez & J. Peter May, editors: *Towards Higher Categories*, Springer, pp. 105–191, doi:10.1007/978-1-4419-1524-5\_4.
- [33] Joachim Lambek (1999): *Diagram Chasing in Ordered Categories with Involution*. *Journal of Pure and Applied Algebra* 143, pp. 293–307, doi:10.1016/S0022-4049(98)00115-7.
- [34] David K. Lewis (1968): *Counterpart Theory and Quantified Modal Logic*. *Journal of Philosophy* 65, pp. 113–26, doi:10.2307/2024555.
- [35] Minghui Ma (2011): *Mathematics of Public Announcements*. In Hans van Ditmarsch, Jérôme Lang & Shier Ju, editors: *Logic, Rationality and Interaction: Third International Workshop, LORI 2011*, Springer, pp. 193–205, doi:10.1007/978-3-642-24130-7\_14.
- [36] Saunders Mac Lane (1998): *Categories for the Working Mathematician*, second edition. Springer-Verlag, doi:10.1007/978-1-4757-4721-8.
- [37] Saunders Mac Lane & Ieke Moerdijk (1992): *Sheaves in Geometry and Logic: A First Introduction to Topos Theory*. Springer-Verlag, doi:10.1007/978-1-4612-0927-0.
- [38] Lawrence S. Moss (1999): *Coalgebraic Logic*. *Annals of Pure and Applied Logic* 96, pp. 277–317, doi:10.1016/S0168-0072(98)00042-6.
- [39] Andrew M. Pitts (2000): *Categorical Logic*. In Samson Abramsky, Dov M. Gabbay & Thomas S. E. Maibaum, editors: *Handbook of Logic in Computer Science, Volume 5: Algebraic and Logical Structures*, Oxford University Press, pp. 39–128.
- [40] S. K. Thomason (1975): *Categories of Frames for Modal Logic*. *Journal of Symbolic Logic* 40, pp. 439–442, doi:10.2307/2272167.